

Case Report

A contribution to flora, life forms and chorology of plants (Case study: South West Ardabil province, Iran)

Mojtaba Akhavan Armaki

Department of Agriculture
Payame Noor University (PNU), Karaj, Iran
Email: mtakhavan@ut.ac.ir

ABSTRACT

Floristic studies are fundamental for the applied sciences such as rangeland management and conservation. Unique ecological and climatic conditions in the South West Ardabil Province (SWA) make it a remarkable habitat for the floristic studies. Plant species were collected from 50 field sites, representing major habitats of winter and rural rangelands. Surveys were conducted during active growth periods in 2012 and 2013. Plant species were identified and their chorology and life form determined through laboratory examinations and by using reference books. Floristic surveys resulted in identification of 534 plant species, belonged to 315 genus and 68 families. Asteraceae, Poaceae and Fabaceae were respectively the most abundant plant families. Among the life forms, Hemicryptophytes, Therophytes, Chamaephytes, Phanerophytes and Geophytes included 45, 32, 9, 7 and 7% of the total species, respectively. Irano-Turanian was the most dominant (65%) Chorotypes. The dominance of Hemicryptophytes and Therophytes, as well as vast distribution of Chamaephytes, can be referred to the simultaneous effects of climate fluctuations and livestock grazing on the flora of winter and rural rangelands in SWA.

Keywords: Rangeland, Floristic composition, South West Ardabil, Iran.

INTRODUCTION

Iran is known as one of the most appealing places for studying plant diversity, as 22% of its 8000 plant species are endemic (Asri 2000). Identifying flora of each region is fundamental for accomplishing other pure and applied researches in biology. Especially in the South West Ardabil (SWA), unique ecological and climatic conditions make it a remarkable habitat for the floristic studies. Winter and rural rangelands comprise a large fraction of SWA area. People in this region are highly dependent on rangelands; they use range plants as sources for food, medicine, livestock production etc. Therefore, identifying floristic list of rural and winter rangelands of SWA is beneficial for protecting the endangered

plants, and for planning a sustainable use of forage, ornamental and medicinal plants. Several reference books (e.g. Rechinger 1967-2010; Boissier 1867-1888; Assadi et al., 2001-2010) provide valuable information on the native and exotic plants of SWA. However, they are mainly adopted from the large-scale studies and do not provide detailed information on the flora of this region. On the other hand several studies have been conducted on the flora of SWA (e.g. Shaad and Sanjari 2011; Aydani 2004; Sobhani et al., 2007) but they only represent information on the flora small areas (usually less than 10000 hectares). Akhaneh (2009) have studied flora of Golestan National Park, in which small parts of

Germeh and Maneh-o-Samalghan counties were also included. Other researchers have studied geographic distribution of plant species for only one family or one genus in Ardabil Provinces, e.g. Compositae (Sobhani and Rajamand, 2007), (Cupressaceae Sadeghhalnejat 2007), and genus of *Bromus* L. (Memariani and Joharchi, 2007). Accordingly, a lack of comprehensive information on the flora rural and winter rangelands of SWA was the most important reason behind this research. The main aim was to do a survey the flora and to identify the major plant phenotype and chorotypes of the winter and rural rangelands in SWA. Result of this study can also be used for the applied researches such as rangeland management and conservation.

MATERIALS AND METHODS

South West Ardabil Province is located between 45° 36' to 26° 39' northern latitude and 5° 45' to 22° 48' eastern longitude. It is located in the northwest of Iran, and comprised of six counties i.e. Kaleybar, Khoda afarin, Ahar, Varzaqan, Jolfa and Marand. Rangelands contain 55.82% of SWA area, which is about 1.72 million hectares, from which 1.31 million hectares are used as winter and rural rangelands (Jankju et al., 2009).

Main bioclimatic region in SWA is Irano-Turanian that is distinguished by hot and dry summers, wet and cold winters. The region is also influenced by cold and dry winds from Siberia (north) and wet and mild climate from the Caspian Sea (west). Therefore the flora of SWA are affected by sever climate fluctuations, because of interaction between the three major climatic regimes. Average yearly rainfall varies from the lowest (<300 mm) in Marand to the highest (650 mm) in the northwest of Kaleybar. Average annual temperature also varies from a maximum (17-18 °C) in Varzaqan to the minimum (8-10 °C) values in Khoda afarin. Many places of SWA winter and rural rangelands confer similar climate, geology and topography conditions. Therefore, site selection was conducted by using geology, topography, soil and seasonal livestock migration

(Kooch) maps. Information provided by the local experts was also implemented. A new site was selected, where the environmental conditions or land use history varied. Accordingly, 50 rangeland sites were selected from which 30 were located in rural and 20 in winter rangelands. Geographic distribution of rural rangeland sites were: 4 in Marand, 9 in Kaleybar, 7 in Khoda afarin, 5 in Varzaqan and 2 in Jolfa. Sites of winter rangelands were scattered as 9 in Marand, 8 in Ahar, 2 in Varzaqan, and one in Kaleybar. Soil textures of most sites were silty loam, but in some places it was clay-loam or silty- clay-loam, pH was about 7-8, soil fertility was low to medium; nitrogen was highly limited whereas potassium was high nearly in all sites (Jankju et al., 2009). Plant species were identified by using some reference books (Akhani 2009; Assadi et al., 2001-2010; Davis 2007; Komarov et al., 1963-1974; Rechinger 1967-2010; Townsend et al., 2006; Tutin et al., 1964-1980); also by botanical identification in the Botanical Research Center of Ferdowsi University of Mashhad. Plant chorotypes were also determined for each plant species, by using several reference books (Akhani 2009; Assadi et al., 2001-2010; Davis 2007; Komarov et al., 1963-1974; Rechinger 1967-2010; Townsend et al., 2006; Tutin et al., 1964-1980; Zohary et al., 1980). Plant phenotypes were determined according to the Raunkiaer's life form specifications (Asri 2000). Plant chorotypes were identified based on Zohary (2004).

RESULTS

Floristic surveys on the winter and rural rangelands of Southern Ardabil Province (NAP) resulted in identification of 534 plant species, belonged to 315 genera, 68 families. Family name, Chorotype and Phenotype of each plant species are compared in the appendix. About 65% of the total plant species in NAP were belonged to the Irano-Turanian Chorotype, whereas Irano-Turanian-Mediterranean, Irano-Turanian-Mediterranean- Euro- Siberian, Irano-Turanian-Euro-Siberian and Pluriregional plant species

respectively contained 11, 10, 6, and 5 percent of all plant species. Less than 3% of total plant species were belonged to the other Chorotypes like cosmopolitans and Subcomopolitans (Fig. 1).




Fig. 1. Plant chorotypes and their relative percentage in flora of winter and rural rangelands in Southern Ardabil province

Plant classification, based on Raunkiaer's life forms revealed Hemicryptophytes as the most abundant (45% of total) species. Therophytes, Chamaephytes, Phanerophytes and Geophytes, only contained 32, 9, 7 and 7 percents of total plant species, respectively (Fig. 2). Among the 68 plant families found in the NAP, Asteraceae, Poaceae and Fabaceae were the most abundant. These families respectively contained 70 (13%), 61 (11%) and 50 (9%) species (Fig. 3).




Fig. 2. Plant life forms and their relative contribution (percent) in flora of winter and rural rangelands in Southern Ardabil Province




Fig. 3. Proportional contribution of plant families in flora of winter and rural rangelands in Southern Ardabil Province; Others: families less than 2% abundance

Nevertheless, there were some shrubs e.g. *Juniperus excelsa*, *Berberis integririma*, *Cotoneaster* spp., *Crataegus* spp., which are typical flora of Armeno-Iranian Phytochorion (Zohary 2004). Armeno-Iranian Phytochorion is one of Irano-Turanian's subdivisions. The presence of plants from other Phytochorions, are mainly due to interaction between three climate conditions; i.e. Mediterranean, Caspian Seas and Euro-Siberian. Hemicryptophytes and Therophytes were the most abundant life form in NAP. Several other studies in Ardabil Province have also reported higher abundance of Therophytes and Hemicryptophytes. Amiri et al. (2008) studied floristic of Tiregan in Hezar Masjed Mts. Memariani et al., (2009) also studied floristic of Fereizi in Chenaran, and both found higher abundance of Hemicryptophytes as compared to other life forms. In studies conducted at other parts of Iran, also higher fractions of Therophytes and Hemicryptophytes have been reported for the rangeland flora. In Genu at Hormozgan Nadjafi et al., (2006) and Garmsar at Semnan (Iranbakhsh et al., 2008) Therophytes were the most abundant life forms. In Fereizi at Azarbaijan, Therophytes and Hemicryptophytes

were commonly the most abundant life forms (Memariani et al., 2009); in Khabr National Park and Rouchoun wildlife refuge (Irannezhad Parizi et al., 2010) and in Meimand (Vakili Shahrbabaki et al., 2011) both in Kerman, and in Kalat highlands of Gonabad in Khorasan Razavi (Vaseghi et al., 2009) Hemicryptophytes were the most abundant plant life forms. Higher frequency of Therophytes and Hemicryptophytes in NAP can be related to their high adaptation to the Mediterranean climate conditions (Zohary 2004). The active growth periods of these life forms are concurrent with the rainy season in late winter and early spring (Tavili et al., 2009). During most of the summer and all winter times, Hemicryptophytes lose their aboveground parts while Therophytes remain as seed. Therefore, these plants avoid summer drought and winter cold stresses (Barbour et al., 1987). Chamaephytes are also Xerophytes plant and are geographically distributed within the whole NAP region. On the other hand, the mesic Phanerophytes and Geophyte species were less abundant in the rural and winter rangelands of NAP; mainly found in the southern slope aspects or at the bank of seasonal water streams. In addition to the climate, intense grazing pressure can also be a determinant factor for the relative abundance and geographic distribution of different life forms (Heithschmidt and Stuth, 1991). Nearly the whole area of NAP is dominated by different species of *Artemisia* L., which contain lots of aromatic materials and essential oils (Cumming and Reid, 2008). Furthermore, thorny shrubs (e.g. *Astragalus* spp., *Acantholimon* spp., and *Acanthophyllum* spp.), poisonous plants (for instance *Euphorbia* spp and *Peganum harmala*) spiny forbs (e.g. *Cirsium* spp., and *Cousinia* spp.) and annual plants (such as *Eremopyron* spp. *Aegilops* spp. and *Bromus tectorum*) dominate vast areas of winter and rural rangelands in NAP. Aromatic and poisonous materials, and spines, or short life cycles are also plant adaptations to livestock grazing (Grime, 2007). In conclusion, winter and rural rangelands of NAP confer a relatively rich floristic

composition, which is a result of plant responses to Mediterranean climate as well as intense livestock grazing. A combination of climate and land use impact has led to dominance of Hemicryptophytes and Therophytes, as well as vast distribution of Chamaephytes in the winter and rural rangelands of NAP. Climate has significant effect on the flora of all habitats in the NAP, whereas human impact is only significant on flora of rural rangeland.

REFERENCES

1. Akhiani, H.,(2009). The illustrated flora of Goleston National Park, Iran. Tehran Univ. Press. Vol. 1, 481pp.
2. Amiri, M.S., Zokaii, M., Ejtehad, H., Mozaffarian, V.,(2008). Study of floristics, life form and chorology of plants in the Tiregan watershed area, Ardabil province, Iran. *Tarbiat Moallem Jour. Sci.*, 8(2): 89-106.
3. Archibold, O.W.,(1996). Ecology of world vegetation. 1st.ed, Chapman and Hall. London. 510 pp.
4. Ardakani, M.R.,(2010). Ecology. Tehran Univ. Press. 340 pp.
5. Asri, U.,(2000). An ecological survey of plant communities of arid regions - Case study: Touran Biosphere Reserve, Semnan Province, Iran. PhD. Thesis, R. & S. unit of Islamic Azad Univ., 302 pp.
6. Assadi, M., Maassoumi, A.A., Khatamsaz, M., Mozaffarian, V., (eds.),(2009). Flora of Iran, Vols. 1-66., Research Institute of Forests and Rangelands Publications. Tehran.
7. Aydani, M.,(2004). Floristic study of Ghoroghe-Darkesh area in northwestern of Khorasan province. M.S. Thesis, R. & S. unit of Islamic Azad Univ., 232 pp.
8. Barbour, M.G., Burk, J.H., Pitts, W.D.,(1987).Terrestrial Plant Ecology., 2nd Edition, The Benjamin/Cummings publishing Company., California, USA.
9. Boissier, P. E.(1867-1888). *Flora Orientalis*, Vols: 1-5. Genevae et Basileae
10. Cumming, J., Reid, N.(2008). Stand-level management of plantations to improve biodiversity values. *Biodiversity and Conservation*. 17(5): 1187-1211.

11. Davis, P. H., (ed.), (2007). Flora of Turkey and the east Aegean Islands., 10 Vols. Edinburgh University Press. UK.
12. Grime, J. P. (2007). Plant Strategies, Vegetation Processes, and Ecosystem Properties. 2nd edition Wiley, Chichester, UK.
13. Heitschmidt, R.K., Stuth, J.W. (1991). Grazing Management: An Ecological Perspective. Timber Press, Portland, Oregon, USA.
14. Iranbakhsh, A.R., Hamdi, S.M. M., Assadi, M. (2008). Flora, life forms and Chorotypes of plants of Garmsar region in Semnan province, Iran, Pajouhesh & Sazandegi, 79: 179-199.
15. Irannezhad Parizi, M. H., Sanei Shariat Panahi, M., Zobeiri, M., Marvi Mohajer, M.R., (2010). Study of floristic and phytogeography in Khabr National Park and Rouchoun wildlife refuge. Iran Jour of Natural Resources. 54(2): 111-128.
16. Jankju, M., Mellati, F., Bozorgmehr, A., (2009). Introduction and investigation of important range plants in the Northern Ardabil Province. Faculty of Natural Resources and Environment, Ferdowsi University of Mashhad, Mashhad, Iran, 235 pp.
17. Komarov, V.L., and. Shishkin, B.K. (1963-1974). Flora of the USSR. Vols. 1-30, Russia.
18. Memariani, F., Joharchi, M. R., (2007). Taxonomy and phytogeography of the genus Bromus L. (Poaceae) in Flora of Azarbaijan., Abstract book of 1st national plant taxonomy conference of Iran. R.I.F.R., P: 32.
19. Memariani F., Joharchi, M.R., Ejtehad, H., Emadzade, Kh., (2009). Contributions to the flora and vegetation of Binalood mountain range, NE Iran: Floristic and chorological studies in Fereizi region. FUIJBS, 1(1): 1-17.
20. Nadjafi Tireh –Shabankareh, K., Jalili, A., Azarbaijan, N., Jamzad Z., Asri, U. (2006). Flora, life forms and chorotypes of plants province (Iran), Pajouhesh & Sazandegi, 69: 50-62.
21. Rechinger, K. H. (1987). Flora Iranica, Volumn 1-178. Graz: Akademische Druck
22. Sadeghalnejat, T. (2007). Revision of the family of Cupressaceae in Khorasan provinces. Abstract book of 1st National Plant Taxonomy Conference of Iran. R.I.F.R., P: 10.
23. Shaad, Q.A., and Sanjari, G. (2011). Plant communities in Ashkhaneh area. R.I.F.R. Press., 72 pp, Tehran, Iran.
24. Sobhani, A., Rajamand, M. 2007. Floristic study of the plant species of Compositae family and investigation of life form, chorotype in Atrak river watershed area in the north Ardabil, Abstract book of 1st national plant taxonomy conference of Iran. R.I.F.R., P: 38.
25. Sobhani, A., Rajamand, M., Naddaf, M. (2007). A Floristic investigation of Shirin Darreh Dam watershed area, NW Bojnourd, North Khorasan province. Abstract book of 1st national plant taxonomy conference of Iran. R.I.F.R. Publication, P: 28.
26. Takhtajan, A. (1986). Floristic Regions of the World, Translated from Russian by T.J. Crovello: Berkeley, University of California Press, 522 pp.
27. Townsend, C.C., Guest, E., Omar, S.A., Alkayat, A.H., (2006). Flora of Iraq. Vols. 1-4. & 8-9., Ministry of Agriculture & Agrarian Reform Republic of Iraq.
28. Tutin, T.G., Heywood, V.H., Burges, N.A., Valentine, D.H., Walters, S.M., Webb, D.A. (1964-1980). Flora European., Vols. 1-5. Cambridge University Press, UK.
29. Vakili Shahrabaki, S.M.A., Atri, M., Assadi, M. (2011). Introduction to the flora, life form and plant geographical distribution of Meimand region in Shahrabak (Kerman), Pajouhesh & Sazandegi. 52: 75-81.
30. Vaseghi, P., Ejtehad, H., Zokaii, M., (2008). Floristic study, life form and chorology of plants in Kalat highlands of Gonabad, Khorasan province, Iran. Tarbiat Moallem Jour. of Sci., 8(1): 75-88.
31. Zohary, M., 2004. Geobotanical Foundations of the Middle East. Vol. 1-2. Gustav Fischer Verlag.
32. Zohary, M., Heyn, C.C., Haller, D., (1980). Conspectus Flora Orientalis., Vols: 1-9, Jerusa

Appendix: Checklist of vascular plant taxa in winter and rural rangelands in southern Ardabil province, Iran.

Life form	Chorotype	Scientific name of plant species in its family
		Aceraceae
Ph.	IT	<i>Acer monspessulanum</i> L. subsp. <i>turcomanicum</i>
		Apiaceae (Umbelliferae)
Th.	IT, M	<i>Ammi majus</i> L.
Th.	IT	<i>Bunium cylindricum</i> (Boiss. & Hohen.) Drude
G.	IT	<i>Bunium persicum</i> (Boiss.) B. Fedtsch.
H.	PI	<i>Conium maculatum</i> L.
H.	IT, ES, M	<i>Falcaria vulgaris</i> Bernh.
H.	IT	<i>Ferula gumosa</i> Boiss.
H.	IT	<i>Ferula ovina</i> (Boiss.) Boiss.
H.	IT, M	<i>Prangos ferulacea</i> (L.) Lindl.
H.	IT	<i>Prangos latiloba</i> Korov.
G.	IT	<i>Elaeosticta allioides</i> (Regel & Schmalh.) Kljuyk. & al.
Th.	IT, M	<i>Scandix stellata</i> Banks & Soland.
Th.	IT, ES, M	<i>Torilis heterophylla</i> Guss.
Th.	PL	<i>Turgenia latifolia</i> (L.) Hoffm.
H.	IT	<i>Zosimia absinthifolia</i> (Vent.) Link.
		Asclepiadaceae
H.	IT	<i>Vincetoxicum pumilum</i> Decne.
		Asteraceae (Compositae)
H.	IT, ES	<i>Achillea biebersteinii</i> Afan
H.	IT, ES	<i>Achillea wilhelmsii</i> C. Koch
G.	IT	<i>Acroptilon repens</i> (L.) DC. subsp. <i>australe</i>
H.	IT, ES	<i>Arctium lappa</i> L.
Ch.	IT, M	<i>Artemisia absinthium</i> L.
Th.	IT, M	<i>Artemisia annua</i> L.
Ch.	IT	<i>Artemisia deserti</i> Krasch.
Ch.	IT	<i>Artemisia kopetdaghensis</i> Krasch., M. Pop. & Lincz. Ex Poljak.
H.	IT, ES, M	<i>Artemisia scoparia</i> Waldst. & Kit
Ch.	IT	<i>Artemisia sieberi</i> Besser
Ch.	IT	<i>Artemisia turanica</i> Krasch.
Th.	IT, ES	<i>Callicephalus nitens</i> (M. B. ex Willd.) C. A. Mey.
Th.	IT, ES	<i>Carduus lanatus</i> L. subsp. <i>turkestanicus</i> (M. Pop.)
Th.	IT, M	<i>Carduus pycnocephalus</i> L.
Th.	IT, M	<i>Carduus transcaspicus</i> Gandog.
Th.	IT	<i>Carthamus oxyacantha</i> M. B.
H.	IT	<i>Centaurea balsamita</i> Lam.
Th.	IT	<i>Centaurea depressa</i> M. B.
H.	IT, ES	<i>Centaurea iberica</i> Trev. ex Spreng.
H.	IT	<i>Centaurea solstitialis</i> L.
G.	IT	<i>Cephalorrhynchus kossinskyi</i> (Krasch.) Kirp.
Th.	IT	<i>Chardinia orientalis</i> (L.) O. Kuntze
H.	IT, ES, M	<i>Chondrilla juncea</i> L.
H.	IT, ES, M	<i>Cichorium intybus</i> L.
H.	PI	<i>Cirsium arvense</i> (L.) Scop. var. <i>incanum</i>
H.	IT	<i>Cirsium bornmulleri</i> Sint. ex. Bornm.
Th.	IT, M	<i>Cnicus benedictus</i> L.
Life form	Chorotype	Scientific name of plant species in its family
H.	IT	<i>Codonocephalum peacockianum</i> Aitch. & Hemsl.
H.	IT	<i>Cousinia arida</i> C. Winkl.
H.	IT	<i>Cousinia congesta</i> Bunge
H.	IT	<i>Cousinia decipiens</i> Boiss. & Buhse
H.	IT	<i>Cousinia komarowii</i> (O. Kuntze) C. Winkl.
H.	IT	<i>Cousinia lasiandra</i> Bunge

H.	IT	<i>Cousinia microcarpa</i> Boiss.
H.	IT	<i>Cousinia turcomanica</i> C. Winkl.
Th.	IT	<i>Crepis turcomanica</i> Krasch.
H.	IT	<i>Gundelia tournefortii</i> L.
H.	IT	<i>Helichrysum graveolens</i> (M. B.) Sweet
H.	IT	<i>Helichrysum oocephalum</i> Boiss.
H.	IT, ES	<i>Heteropappus altaicus</i> (Willd.) Novopokr
G.	IT, ES, M	<i>Inula oculus-christi</i> L.
G.	IT, ES, M	<i>Inula salicina</i> L.
G.	ES	<i>Inula thapsoides</i> (M. B. ex Willd.) Spreng.
G.	IT	<i>Jurinea monocephala</i> Aitch. & Hemsl.
Th.	IT, SS	<i>Koelipinia linearis</i> Pall.
H.	IT, ES, M	<i>Lactuca serriola</i> L.
H.	IT	<i>Lactuca undulata</i> Ledeb.
H.	IT	<i>Launaea acanthodes</i> (Boiss.) O. Kuntze
H.	IT, ES	<i>Leontodon asperrimus</i> (Willd.) Boiss. ex Ball.
H.	IT	<i>Nikitinia leptoclada</i> (Bomm. & Sint.) Iljin
H.	IT	<i>Onopordon heteracanthum</i> C. A. Mey.
H.	IT	<i>Onopordon leptolepis</i> DC.
H.	IT	<i>Picnomon acarna</i> (L.) Cass.
H.	IT	<i>Picris strigosa</i> M. B.
H.	IT, ES, M	<i>Pulicaria dysenterica</i> (L.) Bernh.
H.	IT	<i>Pulicaria gnapholodes</i> (Vent.) Boiss.
H.	IT	<i>Scariola orientalis</i> (Boiss.) Sojak
G.	IT	<i>Scorzonera leptophylla</i> (DC.) Krasch. & Lipsch.
Th.	IT, M	<i>Senecio paulsenii</i> O. Hoffm.
H.	IT	<i>Serratula latifolia</i> Boiss.
H.	IT, M	<i>Sonchus asper</i> (L.) Hill
H.	PI	<i>Tanacetum parthenium</i> (L.) Schultz-Bip.
H.	IT	<i>Tanacetum polycephalum</i> Schultz-Bip.
H.	IT	<i>Taraxacum syriacum</i> Boiss.
H.	IT	<i>Tragopogon graminifolius</i> DC.
Th.	IT	<i>Tripleurospermum disciforme</i> (C. A. Mey.) Schultz-Bip.
G.	IT, ES, M	<i>Tussilago farfara</i> L.
Th.	IT, M	<i>Xanthium spinosum</i> L.
Th.	IT, M	<i>Xanthium strumarium</i> L.
		Berberidaceae
Ph.	IT	<i>Berberis integerrima</i> Bunge
		Boraginaceae
H.	IT, SS	<i>Anchusa italica</i> Retz.
H.	IT	<i>Anchusa ovata</i> Lehm.
Th.	IT	<i>Arnebia decumbens</i> (Vent.) Coss. & Karl
Th.	IT	<i>Arnebia linearifolia</i> DC.
Th.	PI	<i>Asperugo procumbens</i> L.
Life form	Chorotype	Scientific name of plant species in its family
Th.	IT	<i>Buglossoides tenuifolia</i> (L. f.) Johnston.
Th.	IT	<i>Caccinia macranthera</i> (Banks & Soland.) Brand.
H.	IT, M	<i>Echium italicum</i> L.
H.	IT	<i>Heliotropium europaeum</i> L.
H.	IT	<i>Heliotropium lasiocarpum</i> Fisch. & Mey.
Th.	IT	<i>Heterocaryum subsessile</i> Vatke
Th.	IT	<i>Lappula barbata</i> (M. B.) Gurke
Th.	IT	<i>Lappula ceratophora</i> (M. Pop.) M. Pop.
Th.	IT	<i>Lappula microcarpa</i> (Ledeb.) Gurke
Th.	IT	<i>Lappula sinaica</i> (DC.) Ascherson ex Schweinf.
Th.	IT	<i>Nonnea caspica</i> (Willd.) G. Don.
Th.	IT	<i>Nonnea lutea</i> (Desr.) Reichenb.

A contribution to flora, life forms and chorology of plants

H.	IT	<i>Onosma dichroanthum</i> Boiss.
H.	IT	<i>Onosma longilobum</i> Bge.
H.	IT	<i>Paracaryum intermedium</i> (Fresen.) Lipsky
H.	IT	<i>Paracaryum turcomanicum</i> Bornm. et Sint.! ex Bornm.
Th.	IT	<i>Rindera lanata</i> (Lam.) Bunge
Th.	IT	<i>Rochelia disperma</i> (L. f.) C. Koch
H.	IT	<i>Solenanthes circinnatus</i> Ledeb.
H.	IT, M	<i>Solenanthes stamineus</i> (Desf.) Wettst.
H.	IT	<i>Trichodesma incaium</i> (Bge.) A. DC.
		Brassicaceae (Cruciferae)
Th.	IT	<i>Aethionema carneum</i> (Banks & Soland.) B. Fedtsch.
H.	IT, ES, M	<i>Alliaria petiolata</i> (M. B.) Cavara & Grande
Th.	IT	<i>Alyssum dasycarpum</i> Steph. ex Willd.
Th.	IT	<i>Alyssum inflatum</i> Nyarady
Th.	IT	<i>Alyssum linifolium</i> Steph. ex Willd.
Th.	IT, M	<i>Alyssum minus</i> (L.) Rothm.
Th.	IT	<i>Alyssum szowitsianum</i> Fisch. & C. A. Mey.
H.	IT, ES, M	<i>Arabis nova</i> Vill.
H.	IT	<i>Brassica elongata</i> Ehrh.
Th.	IT	<i>Camelina rumelica</i> Velen. subsp. <i>rumelica</i>
Th.	Cos	<i>Capsella bursa-pastoris</i> (L.) Medicus
G.	IT, M	<i>Cardaria draba</i> (L.) Desv.
Th.	IT, SS	<i>Clypeola aspera</i> (Grauer) Turrill
Th.	IT	<i>Clypeola dichotoma</i> Boiss.
Th.	IT	<i>Conringia orientalis</i> (L.) Andrz.
Th.	IT	<i>Conringia perfoliata</i> (C. A. Mey.) Busch
H.	IT	<i>Crambe kotschyana</i> Boiss.
Th.	IT, ES, M	<i>Descurainia Sophia</i> (L.) Schur
Th.	IT	<i>Eruca sativa</i> Lam.
H.	IT	<i>Erysimum aitchisonii</i> O. E. Schulz
H.	IT	<i>Erysimum ischnostylum</i> Freyn & Sint.
Th.	IT, M	<i>Euclidium syriacum</i> (L.) R. Br.
Th.	IT, M	<i>Euclidium tenuissimum</i> (Pall.) B. Fedtsch.
H.	IT	<i>Fibigia suffruticosa</i> (Vent.) Sweet
Th.	IT, M	<i>Goldbachia laevigata</i> (M. B.) DC.
H.	IT	<i>Graellsia integrifolia</i> (Rech. f.) Rech. f.
H.	IT, ES, M	<i>Isatis leuconeura</i> Boiss. & Buhse
Life form	Chorotype	Scientific name of plant species in its family
H.	IT	<i>Lepidium latifolium</i> L.
H.	IT	<i>Leptaleum filifolium</i> (Willd.) DC.
Th.	IT	<i>Malcolmia africana</i> (L.) R. Br.
Th.	IT	<i>Malcolmia strigosa</i> Boiss.
Th.	IT	<i>Matthiola chenopodifolia</i> Fisch. & C. A. Mey.
Th.	IT	<i>Matthiola farinosa</i> Bge. ex Boiss.
H.	IT	<i>Peltaria turkmena</i> Lipsky
Th.	IT, ES, M	<i>Rapistrum rugosum</i> (L.) All.
Th.	IT, ES, M	<i>Sisymbrium altissimum</i> L.
Th.	IT	<i>Thlaspi kotschyana</i> Boiss. & Hohen.
Th.	IT, ES, M	<i>Thlaspi perfoliatum</i> L.
		Caesalpinaceae
Ph.	IT, M	<i>Cercis griffithii</i> Boiss.
		Campanulaceae
H.	IT, ES	<i>Campanula glomerata</i> L.
		Capparidaceae
Ph.	IT, SS, M	<i>Capparis spinosa</i> L.
H.	IT	<i>Cleome heratensis</i> Bunge & Boiss.
H.	IT	<i>Cleome Khorasanica</i> Bunge & Bien. ex Boiss.

		Caprifoliaceae
Ph.	IT, ES	<i>Lonicera floribunda</i> Boiss. & Buhse
Ph.	IT, M	<i>Lonicera nummularifolia</i> Jaub. & Spach
H.	IT, M	<i>Sambucus ebulus</i> L.
		Caryophyllaceae
Ch.	IT	<i>Acanthophyllum bracteatum</i> Boiss.
Ch.	IT	<i>Acanthophyllum glandulosum</i> Bunge ex Boiss.
Ch.	IT	<i>Acanthophyllum microcephalum</i> Boiss.
H.	IT	<i>Buffonia oliveriana</i> Ser.
H.	IT	<i>Buffonia sintenisii</i> Freyn
H.	IT	<i>Dianthus crinitus</i> Sm. subsp. <i>turcomanicus</i> (Schischk.) Rech. f.
H.	IT	<i>Dianthus orientalis</i> Adams subsp. <i>stenocalyx</i> (Boiss.) Rech. f.
Ch.	IT	<i>Gypsophila aretioides</i> Boiss.
H.	IT	<i>Gypsophila bicolor</i> (Freyn & Sint.) Grossh.
Th.	IT	<i>Holosteum glutinosum</i> (M. B.) Fisch. & C. A. Mey.
H.	IT	<i>Minuartia meyeri</i> (Boiss.) Bomm.
Ch.	IT	<i>Saponaria bodeana</i> Boiss.
H.	IT	<i>Silene bupleuroides</i> L.
H.	IT, ES	<i>Silene conoidea</i> L.
H.	IT, M	<i>Vaccaria oxyodonta</i> Boiss.
		Chenopodiaceae
Th.	IT	<i>Anabasis aphylla</i> L.
Ch.	IT	<i>Atriplex canescens</i> James
Th.	IT	<i>Atriplex flabellum</i> Bge.
Th.	IT	<i>Atriplex moneta</i> Bge.
Th.	IT	<i>Atriplex tatarica</i> L.
Ch.	IT	<i>Atriplex verrucifera</i> M. B.
Th.	IT	<i>Ceratocarpus arenarius</i> L.
Th.	IT	<i>Chenopodium album</i> L.
Th.	IT, M	<i>Chenopodium botrys</i> L.
Th.	IT	<i>Chenopodium foliosum</i> (Moench) Asch.
Life form	Chorotype	Scientific name of plant species in its family
Th.	IT	<i>Gamanthus gamocarpus</i> (Moq.) Bge.
Th.	IT	<i>Halimocnemis pilifera</i> Moq.
Ch.	IT	<i>Halocnemum strobilaceum</i> M. B.
Ch.	IT	<i>Halothamnus subaphyllus</i> (C. A. Mey.) Botsch.
Ch.	IT	<i>Holostachys caspica</i> (Pall.) C. A. Mey.
H.	IT	<i>Kochia prostrata</i> (L.) Schrad.
Th.	IT	<i>Kochia scoparia</i> (L.) Schrad.
Ch.	IT	<i>Krascheninnikowia ceratoides</i> (L.) Gueldenst.
H.	IT	<i>Noaea mucronata</i> (Forsk.) Asch. et Schweinf.
Ch.	IT	<i>Salsola arbusculiformis</i> Drob.
Ch.	IT	<i>Salsola aucheri</i> (Moq.) Bge.
Ch.	IT	<i>Salsola dendroides</i> Pall.
Th.	IT	<i>Salsola gossypina</i> Bge.
Th.	IT	<i>Salsola kali</i> L.
Th.	IT	<i>Salsola nitraria</i> Pall.
Ch.	IT	<i>Salsola orientalis</i> S. G. Gmel.
Th.	IT	<i>Salsola sclerantha</i> C. A. Mey.
Th.	IT	<i>Seidlitzia florida</i> (M. B.) Boiss.
Th.	IT, M	<i>Seidlitzia rosmarinus</i> (Ehrh.) Bge.
Th.	IT	<i>Spinacia oleracea</i> L.
Th.	IT	<i>Spinacia turkestanica</i> Iljin
Th.	IT	<i>Suaeda altissima</i> (L.) Pall.
Th.	IT	<i>Suaeda microphylla</i> Pall.
Th.	IT	<i>Suaeda microsperma</i> (C. A. Mey.) Fenzl
		Cistaceae

A contribution to flora, life forms and chorology of plants

Ch.	IT, ES, M	<i>Fumana procumbens</i> (Dun.) Gren. & Godron
Th.	IT, SS, M	<i>Helianthemum ledifolium</i> (L.) Miller
		Convolvulaceae
G.	Sco	<i>Convolvulus arvensis</i> L.
H.	IT	<i>Convolvulus commutatus</i> Boiss.
H.	IT, M	<i>Convolvulus dorycnium</i> L.
H.	H.	<i>Convolvulus fruticosus</i> Pall.
H.	H.	<i>Convolvulus pseudocantabrica</i> Schrenk
		Crassulaceae
H.	IT	<i>Rosularia paniculata</i> (Regel & Schmalh.) Berger
		Cuscutaceae
Th.	IT	<i>Cuscuta approximata</i> Babingt
Th.	IT	<i>Cuscuta monogyna</i> Vahl
		Cupressaceae
Ph.	IT	<i>Juniperus excelsa</i> M. B.
		Cyperaceae
G.	IT, M	<i>Carex physodes</i> M. B.
G.	IT, M	<i>Cyperus longus</i> L.
		Dipsacaceae
H.	IT	<i>Cephalaria kotschyi</i> Boiss. & Hohen.
H.	IT	<i>Cephalaria microcephala</i> Boiss.
H.	IT	<i>Dipsacus laciniatus</i> L.
H.	ES	<i>Dipsacus strigosus</i> Willd. ex Roemer & Schultes
Th.	IT	<i>Scabiosa olivieri</i> Coult.
Th.	IT	<i>Scabiosa rotata</i> M. B.
Life form	Chorotype	Scientific name of plant species in its family
		Elaeagnaceae
Ph.	IT	<i>Elaeagnus angustifolia</i> L.
		Ephedraceae
Ph.	IT, ES, M	<i>Ephedra major</i> Host
		Euphorbiaceae
Th.	IT	<i>Chrozophora hierosolymitana</i> Spreng.
H.	IT	<i>Euphorbia aucheri</i> Boiss.
Th.	IT	<i>Euphorbia bungei</i> Boiss.
Th.	IT	<i>Euphorbia falcata</i> L.
Th.	IT	<i>Euphorbia inderiensis</i> Less. ex Kar. & Kir.
Th.	IT	<i>Euphorbia szovitsii</i> Fisch. & C. A. Mey.
		Fumariaceae
Th.	IT, ES, M	<i>Fumaria asepala</i> Boiss.
Th.	IT	<i>Fumaria indica</i> (Hauskn.) Pugsley
Th.	IT	<i>Fumaria vaillantii</i> Loisel.
		Fabaceae
H.	IT	<i>Alhagi persarum</i> Boiss. & Buhse
H.	IT	<i>Astragalus ankylotus</i> Fisch. & C. A. Mey.
Th.	IT	<i>Astragalus bakaliensis</i> Bunge
H.	IT	<i>Astragalus brevidens</i> Freyn & Sint.
H.	IT	<i>Astragalus bassineri</i> Boiss. & Hausk. ex Boiss.
Th.	IT	<i>Astragalus campylorrhynchus</i> Fisch. & Mey.
H.	IT	<i>Astragalus cercidophacos</i> Podlech & Maassoumi
H.	IT	<i>Astragalus chrysostachys</i> Boiss.
H.	IT	<i>Astragalus citrinus</i> Bge. subsp. <i>citrinus</i>
H.	IT	<i>Astragalus curvipes</i> Trautv.
H.	IT	<i>Astragalus hermannii</i> Freitag & Podl.
H.	IT	<i>Astragalus khoshjailensis</i> Sirj. & Rech. f.
H.	IT	<i>Astragalus kopetdaghi</i> Boriss. var. <i>orientikoptdghensis</i>
H.	IT	<i>Astragalus siversianus</i> Pall.
Th.	IT	<i>Astragalus maymanensis</i> Podl.

H.	IT	<i>Astragalus mazandranus</i> Bge.
H.	IT	<i>Astragalus pseudoszovitsii</i> Sirj. & Rech. f.
Th.	IT	<i>Astragalus raddei</i> N. Basil.
H.	IT	<i>Astragalus rawlinsianus</i> Aitch. & Baker
Ch.	IT	<i>Astragalus siliquosus</i> Boiss. subsp. <i>siliquosus</i>
H.	IT	<i>Astragalus squarosus</i> Bunge
Ph.	IT, ES	<i>Astragalus suluklensis</i> Freyn & Sint.
Ph.	IT	<i>Colutea buhsei</i> (Boiss.) Shap.
H.	IT, ES	<i>Colutea porphyrogramma</i> Rech. f.
G.	IT, ES, M	<i>Coronilla varia</i> L. subsp. <i>varia</i>
H.	IT	<i>Glycyrrhiza glabra</i> L.
H.	IT	<i>Hedysarum kopetdaghi</i> Boriss.
H.	IT	<i>Hedysarum wrightianum</i> Aitch. & Baker
Th.	IT, M	<i>Lathyrus cicera</i> L.
Th.	IT, ES, M	<i>Lathyrus sativus</i> L.
H.	PI	<i>Lotus corniculatus</i> L.
H.	PI	<i>Medicago lupulina</i> L.
H.	IT, ES, M	<i>Medicago sativa</i> L.
Life form	Chorotype	Scientific name of plant species in its family
H.	IT, ES, M	<i>Melilotus albus</i> Desr.
H.	IT, ES, M	<i>Melilotus officinalis</i> (L.) Desr.
H.	IT	<i>Meristotropis xanthioides</i> Vassilez.
Th.	IT	<i>Onobrychis aucheri</i> Boiss.
Ch.	IT	<i>Onobrychis cornuta</i> (L.) Desv.
H.	IT	<i>Onobrychis sintenisii</i> Bornm.
Ch.	IT	<i>Ononis spinosa</i> L.
Ch.	IT	<i>Prosopis farcta</i> (Banks & Soland.) Macbr.
H.	IT	<i>Sophora alopecuroides</i> L.
H.	IT	<i>Sophora pachycarpa</i> C. A. Mey.
H.	IT, ES, M	<i>Trifolium pratense</i> L.
H.	IT, ES, M	<i>Trifolium repens</i> L.
Th.	IT	<i>Vicia monantha</i> Retz.
Th.	IT, ES, M	<i>Vicia sativa</i> L.
Th.	IT	<i>Vicia subvillosa</i> (Ledeb.) Boiss.
H.	Th.	<i>Vicia variabilis</i> Freyn & Sint.
Th.	IT, ES, M	<i>Vicia villosa</i> Roth
		Gentianaceae
Th.	IT, ES, M	<i>Centaurium pulchellum</i> (Swartz) Druce
		Geraniaceae
G.	IT	<i>Biebersteinia multifida</i> DC.
Th.	IT, ES, M	<i>Erodium cicutarium</i> (L.) L Her.
Th.	IT, ES, M	<i>Geranium rotundifolium</i> L.
		Hypericaceae
H.	IT, M	<i>Hypericum perforatum</i> L.
H.	IT	<i>Hypericum scabrum</i> L.
		Iridaceae
G.	IT	<i>Iris fosterana</i> Aitch. & Baker
G.	IT	<i>Iris kopetdagensis</i> (Vved.) Mathew & Wendelbo
G.	IT	<i>Iris songarica</i> Schrenk
		Juglandaceae
Ph.	IT, ES, M	<i>Juglans regia</i> L.
		Juncaceae
G.	PI	<i>Juncus fontanesii</i> Gay
		Lamiaceae (Labiatae)
H.	IT, ES	<i>Ajuga comata</i> Stapf
Ch.	IT	<i>Dracocephalum subcapitatum</i> (O. Kuntze) Lipsky
H.	IT	<i>Eremostachys boissieriana</i> Regel

H.	IT	<i>Eremostachys labiosiformis</i> (M. Pop.) Knorring
Ch.	IT	<i>Hymenocrater calycinus</i> (Boiss.) Benth.
H.	IT	<i>Hymenocrater elegans</i> Bge.
Ch.	IT	<i>Lagochilus aucheri</i> Boiss.
Ch.	IT	<i>Lagochilus cabulicus</i> Benth.
Th.	IT, M	<i>Lallemantia iberica</i> (Stev.) Fisch. & C. A. Mey.
H.	IT, M	<i>Marrubium anisodon</i> C. Koch
H.	Pl	<i>Marrubium vulgare</i> L.
H.	IT	<i>Mentha longifolia</i> (L.) Hudson
H.	IT, ES, M	<i>Nepeta cataria</i> L.
H.	IT	<i>Nepeta glomerulosa</i> Boiss.
Th.	IT	<i>Nepeta saccharata</i> Bunge
Life form	Chorotype	Scientific name of plant species in its family
Ch.	IT	<i>Perovskia abrotanoides</i> Karel.
H.	IT	<i>Phlomis cancellata</i> Bunge
H.	IT	<i>Phlomis herba-venti</i> L.
H.	IT	<i>Salvia atropatana</i> Bge.
H.	IT	<i>Salvia ceratophylla</i> L.
H.	IT	<i>Salvia chloroleuca</i> Rech. f. & Aell.
H.	IT, M	<i>Salvia sclarea</i> L.
H.	IT, M	<i>Salvia virgata</i> Jacq.
Ch.	IT	<i>Scutellaria litwinowii</i> Bornm. & Sint. ex Bornm.
Ch.	IT	<i>Scutellaria pinnatifida</i> A. Hamilt.
H.	IT, ES	<i>Stachys byzanthina</i> C. Koch
G.	IT	<i>Stachys lavandulifolia</i> Vahl
H.	IT	<i>Stachys pubescens</i> Ten.
H.	IT	<i>Stachys setifera</i> C. A. Mey.
Ch.	IT	<i>Stachys subaphylla</i> Rech. f.
Ch.	IT	<i>Stachys trinervis</i> Aitch. & Hem
H.	IT	<i>Stachys turcomanica</i> Trautv.
H.	IT, ES, M	<i>Teucrium chamaedrys</i> L.
H.	IT, M	<i>Teucrium polium</i> L.
Ch.	IT	<i>Thymus transcaspicus</i> Klokov
Ch.	IT	<i>Thymus trautvetteri</i> Klokov & Desj.-Shost
Ch.	IT	<i>Ziziphora clinopodioides</i> Lam.
Th.	IT	<i>Ziziphora tenuir</i> L.
		Linaceae
Th.	IT, M	<i>Linum glaucum</i> Boiss. & Nöe
Th.	-	<i>Linum</i> sp.
		Liliaceae
G.	IT	<i>Colchicum robustum</i> (Bunge) Stef.
G.	IT	<i>Eremurus olgae</i> Regel
G.	IT	<i>Eremurus stenophyllus</i> (Boiss. & Buhse) Baker
G.	IT	<i>Fritillaria gibbosa</i> Boiss.
G.	IT, ES, M	<i>Muscari neglectum</i> Guss.
G.	IT	<i>Tulipa montana</i> Lindl.
		Lythraceae
H.	IT	<i>Lythrum hyssopifolium</i> L.
H.	Pl	<i>Lythrum salicaria</i> L.
		Malvaceae
H.	IT, M	<i>Althaea cannabina</i> L.
Th.	IT, M	<i>Althaea hirsuta</i> L.
H.	IT	<i>Althaea officinalis</i> L.
Th.	IT	<i>Hibiscus trionum</i> L.
H.	IT, ES, M	<i>Malva neglecta</i> Wallr.
H.	IT	<i>Malva sylvestris</i> L.
		Moraceae

A contribution to flora, life forms and chorology of plants

Ph.	IT, M	<i>Ficus carica</i> L.
Ph.	IT, M	<i>Ficus rupestris</i> (Hauskn. ex Boiss.) Azizian
		Oleaceae
Ph.	IT, ES, M	<i>Jasminum fruticans</i> L.
		Onagraceae
G.	Pl	<i>Epilobium hirsutum</i> L.
Life form	Chorotype	Scientific name of plant species in its family
G.	Pl	<i>Epilobium minutiflorum</i> Hauskn.
		Oxalidaceae
H.	Pl	<i>Oxalis corniculata</i> L.
		Papaveraceae
Th.	IT, M	<i>Glaucium paucilobum</i> Freyn
Th.	IT, M	<i>Hypocoum pendulum</i> L.
Th.	IT	<i>Papaver dubium</i> L.
Th.	IT	<i>Roemeria refracta</i> DC.
		Plantaginaceae
H.	IT, M	<i>Plantago lanceolata</i> L.
H.	Sco	<i>Plantago major</i> L.
		Platanaceae
Ph.	IT, M	<i>Platanus orientalis</i> L.
		Plumbaginaceae
Ch.	IT	<i>Acantholimon avenaceum</i> Bunge
Ch.	IT	<i>Acantholimon erinaceum</i> (Jaub. & Spach) Lincz.
Ch.	IT	<i>Acantholimon pterostegium</i> Bunge
H.	IT	<i>Limonium gmelini</i> (Willd.) O. Kuntze
H.	IT	<i>Limonium reniforme</i> (Girard) Lincz.
H.	IT, ES, M	<i>Plumbago europaea</i> L.
		Podophyllaceae
G.	IT	<i>Bongardia chrysogonum</i> (L.) Boiss.
G.	IT	<i>Leontice leontopetalum</i> L.
		Primulaceae
Th.	Pl	<i>Anagallis arvensis</i> L. var. <i>coerulea</i> L.
Ch.	IT	<i>Dionysia tapetodes</i> Bge.
		Poaceae (Gramineae)
Th.	IT	<i>Aegilops tauschii</i> Cosson
Th.	IT	<i>Aegilops triuncialis</i> L.
H.	IT	<i>Aeluropus littoralis</i> (Gouan) Parl.
H.	IT	<i>Agropyron intermedium</i> (Host) P. Beauv.
H.	IT	<i>Agropyron pectiniforme</i> Roemer & Schultes
H.	IT	<i>Agropyron podperae</i> Nab.
H.	IT	<i>Agropyron trichophorum</i> (Link) Richter
H.	IT	<i>Agrostis olympica</i> (Boiss.) Bor
H.	IT	<i>Agrostis stolonifera</i> L.
H.	IT, M	<i>Alopecurus apiatus</i> Ovcz.
H.	IT	<i>Arrhenatherum kotschyi</i> Boiss.
Th.	Pl	<i>Avena eriantha</i> Durieu
Th.	Pl	<i>Avena fatua</i> L.
Th.	IT	<i>Boissiera squarrosa</i> (Banks & Soland.) Nevski
H.	Pl	<i>Bothriochloa ischaemum</i> (L.) Keng
H.	Pl	<i>Brachypodium sylvaticum</i> (Huds.) P. Beauv.
Th.	IT	<i>Bromus briziformis</i> Fisch. & C. A. Mey.
Th.	IT	<i>Bromus danthoniae</i> Trin.
H.	IT	<i>Bromus kopetdaghensis</i> Drobov
Th.	IT, ES, M	<i>Bromus scoparius</i> L.
Th.	IT, ES, M	<i>Bromus sterilis</i> L.
Th.	IT, SS, M	<i>Bromus tectorum</i> L.
Life form	Chorotype	Scientific name of plant species in its family

A contribution to flora, life forms and chorology of plants

H.	IT	<i>Calamagrostis epigejos</i> (L.) Roth
H.	IT	<i>Calamagrostis pseudophragmites</i> (Hall. f.) Koel.
G.	IT	<i>Cynodon dactylon</i> (L.) Pers.
H.	IT	<i>Dactylis glomerata</i> L.
Th.	IT, M	<i>Digitaria sanguinalis</i> (L.) Scop.
Th.	IT, M	<i>Echinaria capitata</i> (L.) Desf
Th.	IT	<i>Echinochloa crus-galli</i> (L.) P. Beauv.
H.	IT	<i>Elymus baldschuanicus</i> Roshev.
Th.	IT	<i>Eremopyrum bonaepartis</i> (Spreng.) Nevski
Th.	IT	<i>Eremopyrum distans</i> (C. Koch) Nevski
H.	IT, ES	<i>Festuca arundinacea</i> Schreb.
H.	IT, ES	<i>Festuca ovina</i> L.
H.	IT, ES	<i>Festuca sulcata</i> (Hack.) Beck
H.	IT, M	<i>Hordeum bulbosum</i> L.
H.	IT, ES, M	<i>Hordeum violaceum</i> Boiss. et Huet
H.	IT	<i>Lolium persicum</i> Boiss. & Hohen. ex Boiss.
Th.	IT, ES, M	<i>Lolium rigidum</i> Guadin
H.	IT, ES, M	<i>Melica ciliata</i> L.
H.	IT	<i>Melica persica</i> Kunth
Th.	IT	<i>Nardurus subulatus</i> (Banks & Soland.) Bor
H.	IT, SS	<i>Pennisetum orientale</i> L. C. Rich.
Th.	IT, M	<i>Phalaris minor</i> Retz.
H.	IT	<i>Phleum paniculatum</i> Hudson
H.	IT, ES, M	<i>Phleum pratense</i> L.
G.	IT, ES, M, SS	<i>Phragmites australis</i> (Cav.) Trin. ex Steud.
H.	IT, ES, M	<i>Poa bulbosa</i> L.
Th.	PI	<i>Polypogon fugax</i> Nees ex Steud.
Th.	PI	<i>Setaria verticillata</i> (L.) P. Beauv.
Th.	PI	<i>Setaria viridis</i> (L.) P. Beauv.
G.	PI	<i>Sorghum halepense</i> (L.) Pers.
H.	IT	<i>Stipa barbata</i> Desf.
H.	IT	<i>Stipa caucasica</i> Schmalh.
H.	IT	<i>Stipa hohenakeriana</i> Trin. & Rupr
H.	IT	<i>Stipa parviflora</i> Desf.
H.	IT	<i>Stipa pennata</i> L.
H.	IT	<i>Stipagrostis pulmosa</i> (L.) Munro ex. T. Anders
Th.	IT	<i>Taeniatherum crinitum</i> (Schreb.) Nevski
		Polygonaceae
Ch.	IT	<i>Atraphaxis spinosa</i> L.
H.	IT, M	<i>Polygonum hydropiper</i> L.
H.	IT	<i>Polygonum hyrcanicum</i> Rech. f.
H.	IT	<i>Polygonum patulum</i> M. B.
Ch.	IT	<i>Pteropyrum aucheri</i> Jaub. & Spach
H.	IT, M	<i>Rumex thjanschanicus</i> Los.
IT	IT	Ranunculaceae
Th.	IT	<i>Adonis flammea</i> Jacq.
Th.	IT	<i>Anemone biflora</i> DC.
Th.	IT	<i>Clematis orientalis</i> L.
Th.	IT	<i>Consolida regulosa</i> (Boiss.) Schrod.
Life form	Chorotype	Scientific name of plant species in its family
Th.		
	IT	<i>Delphinium turkmenum</i> Lipsky
Th.	IT	<i>Nigella nigellastrum</i> (L.) Willk
Th.	IT, ES, M	<i>Ranunculus arvensis</i> L.
G.	IT	<i>Ranunculus brutius</i> Ten.
G.	IT	<i>Ranunculus cicutarius</i> Schlect.

H.	IT	<i>Thalictrum sultanabadense</i> Stapf
		Resedaceae
H.	IT	<i>Reseda aucheri</i> Boiss.
H.	IT, ES, M	<i>Reseda lutea</i> L.
H.	IT, ES, M	<i>Reseda luteola</i> L.
		Rhamnaceae
Ph.	IT, ES, M	<i>Paliurus spina-christi</i> Miller
IT	IT	Rosaceae
H.	IT, ES, M	<i>Agrimonia eupatoria</i> L.
Ph.	IT	<i>Amygdalus scoparia</i> Spach
Ph.	IT	<i>Cerasus microcarpa</i> (Mey.) Boiss.
Ph.	IT	<i>Cerasus pseudoprostrata</i> Pojark.
Ph.	IT, ES	<i>Cotoneaster nummularia</i> Fisch. & C. A. Mey.
Ph.	IT	<i>Cotoneaster ovata</i> Pojark.
Ph.	IT	<i>Crataegus assadii</i> Khatamsaz
Ph.	IT	<i>Crataegus pseudoheterophylla</i> Pojark.
Ph.	IT	<i>Crataegus turkestanica</i> Pojark.
H.	IT	<i>Geum kokanicum</i> Regel & Schmalh. ex Regel
H.	IT, ES	<i>Geum urbanum</i> L.
H.	IT, ES	<i>Potentilla inclinata</i> Vill.
H.	IT, ES	<i>Potentilla nuda</i> Boiss.
H.	IT, ES	<i>Potentilla recta</i> L.
H.	IT, ES	<i>Potentilla reptans</i> L.
Ph.	IT, ES	<i>Prunus divaricata</i> Ledeb.
Ph.	IT	<i>Rosa baggeriana</i> Schrenk
Ph.	IT, ES, M	<i>Rosa canina</i> L.
Ph.	IT	<i>Rosa hemisphaerica</i> J. Herrmann
Ph.	IT	<i>Rosa iberica</i> Stev.
Ch.	IT	<i>Rosa persica</i> Mich. ex Juss.
Ph.	IT	<i>Rubus sanctus</i> Shreb.
H.	IT, ES, M	<i>Sanguisorba minor</i> Scop. subsp. <i>muricata</i> (Spach)
Briq.		
IT	IT	Rubiaceae
Th.	IT, ES, M	<i>Asperula arvensis</i> L.
Ch.	IT	<i>Asperula glomerata</i> (M. B.) Griseb.
H.	IT, M	<i>Callipeltis cucullaria</i> (L.) Stev.
H.	IT	<i>Crucianella chlorostachys</i> Fisch. & C. A. Mey.
H.	IT	<i>Crucianella gilanicum</i> Trin.
Th.	IT	<i>Gaillonia Bruguierii</i> A. Rich
Th.	IT, M	<i>Galium humifusum</i> M. B.
Th.	IT, M	<i>Galium setaceum</i> Lam.
Th.	IT	<i>Galium tinctorium</i> L.
H.	PI	<i>Galium verum</i> L. subsp. <i>verum</i>
Ch.	IT	<i>Rubia florida</i> Boiss.
Life form	Chorotype	Scientific name of plant species in its family
		Rutaceae
H.	IT	<i>Haplophyllum perforatum</i> (M. B.) Kar. & Kir.
		Salicaceae
Ph.	IT	<i>Salix alba</i> L.
		Santalaceae
G.	IT	<i>Thesium kotschyianum</i> Boiss.
IT	IT	Scrophulariaceae
H.	IT	<i>Leptorhabdos parviflora</i> (Benth.) Benth.
H.	IT, ES	<i>Linaria lineolata</i> Boiss.
H.	IT, ES	<i>Linaria pyramidata</i> (Lam.) Spreng.
H.	IT	<i>Scrophularia striata</i> Boiss.

A contribution to flora, life forms and chorology of plants

H.	IT, ES	<i>Verbascum cheiranthifolium</i> Boiss.
H.	IT	<i>Veronica anagallis-aquatica</i> L.
H.	IT	<i>Veronica Khorasanica</i> Czernjak.
		Simaroubaceae
Ph.	IT	<i>Ailanthus altissima</i> (Mill.) Swingle
		Solanaceae
H.	IT	<i>Hyoscyamus squarrosus</i> Griff.
Ph.	IT	<i>Lycium ruthenicum</i> Murray
Th.	Cos	<i>Solanum nigrum</i> L.
Th.	IT	<i>Solanum olgae</i> Pojark.
		Tamaricaceae
Th.	-	<i>Tamarix sp.</i>
Th.	IT	<i>Reaumuria alternifolia</i> (Labill.) Brihen
		Thymelaeaceae
H.	IT	<i>Dendrostellera lessertii</i> (Wikstr.) Van Tigeh
H.	IT	<i>Stelleropsis antoninae</i> Pobedimova
Th.	IT, ES, M	<i>Thymelaea passerina</i> (L.) Cosson & Germ.
		Ulmaceae
Ph.	ES, M	<i>Celtis caucasica</i> Willd.
		Urticaceae
Ch.	IT, ES, M	<i>Parietaria judaica</i> L.
H.	Sco	<i>Urtica dioica</i> L.
		Valerianaceae
G.	IT	<i>Valeriana ficariifolia</i> Boiss.
H.	IT	<i>Valeriana sisymbiifolia</i> Vahl
Th.	IT, M	<i>Valerianella cymbicarpa</i> C. A. Mey.
Th.	IT	<i>Valerianella oxyrrhyncha</i> Fisch. & C. A. Mey.
H.	IT	<i>Valerianella szowitsiana</i> Fisch. & C. A. Mey.
		Verbenaceae
H.	IT	<i>Verbena officinalis</i> L.
Ph.	IT	<i>Vitex pseudo-negundo</i> (Hauskn.) Hand-Mzt.
		Violaceae
Th.	IT	<i>Viola occulta</i> Lehmann
Th.	IT	<i>Viola odorata</i> L.
		Vitaceae
Ph.	IT	<i>Vitis vinifera</i> L.
Life form	Chorotype	Scientific name of plant species in its family
		Zygophyllaceae
H.	IT, SS, M	<i>Peganum harmala</i> L.
Th.	IT, M	<i>Tribulus terrestris</i> L.
Ph.	IT	<i>Zygophyllum fabago</i> L.