

Research Article**Dry Eye Syndrome and Associated Risk Factors among
Computer Users in Karachi, Pakistan**

¹Hafiza Amna Amin, ²Wasim Shahzad, ³Yasin Khan
and ⁴Khurram Shahzad

¹Allied Hospital, Faisalabad, Pakistan

²DHQ Hospital, Faisalabad, Pakistan

³BHU Alizo, Bajaur, Pakistan

⁴Hamdard University, Karachi, Pakistan

ABSTRACT

Objective: To assess the frequency of presence of dry eyes and its associated risk factors among employees working in call centers and software houses in Karachi.

Methods: A cross-sectional study was done during July to October 2014 at Hamdard College of Medicine and Dentistry, Karachi. By convenient sampling technique, data was collected from employees aged 20 to 40 years of call centers and software houses. Demographic information was recorded one-designed questionnaire and Schemer's test strips were used to measure production of tears. De-scripture analysis was done for categorical variables using SPSS version 20. Cross-tabulation of different variables was done with dryness of eyes experienced. The frequency of various eye complaints was determined and Chi square test was used to observe the association with Schemer's test. P-value<0.05 was considered as significant.

Results: Out of 150 employees 42 (28%) were found to have dry eyes. Mean age was 27 ± 3.5 years. The majority of employees 117 (78%) were males and 77 (51.3%) were using corrective spectacles for refractive errors. The most common eye complaints among the computer users were burning of eyes (44%), redness of eye (41.3%), dryness of nose (36%), dryness of mouth (34.7%), discharge from eye (10.7%) and foreign body sensation (10%). Statistically significant ($p<0.05$) relationship has been observed between presence of eye burning and positive Schemer's test.

Conclusion: This study indicated that 28% of the individuals had dry eye, along with other symptoms of eyes among computer users in call centers and software houses.

Keywords: Dry eye syndrome, asthenia, eyestrain, computer, call centers.

INTRODUCTION

Dry eye syndrome (DES) or kerato conjunctivitis sicca (KCS), is a multifactorial eye disease most common in females and old patients. It is caused by either decreased tear production or increased tear film evaporation normally functioning. 2 tear film is required to maintain clarity of vision and ocular health. The tear film consists of three layers from inside to outside; mucinous layer, aqueous layer, and sebaceous layer which serves to provide the ocular surface comfort,

mechanical, environmental, and immune protection, maintain epithelial cellular health, and provide a smooth and very powerful refracting surface for clear vision. Abnormality of any one of the three layers of tears produces an unstable tear film, resulting in symptoms of KCS. 4 It has been reported in the past that older women suffer more from dry eye syndrome than males. A research conducted among men in the 5, 6 United States concluded that the risk of dry eye

increases with age, presence of hypertension, benign prostatic hyperplasia and use of antidepressants 7 Another study conducted on elderly population in Australia reported the incidences of symptoms of dry eyes are more common in older females 8 Another study in a developing country revealed prevalence of 27.5% of dry eyes symptoms with pterygium; an independent risk factor A 9 cross-sectional hospital-based study done in India reported prevalence of 18.4% with relative increase in symptoms due to excessive wind, sunlight exposure, smoking, air pollution and drug usage 10 In this advanced era of technology, throughout the world the use of computers has become a necessity in every working environment. This has significantly increased the impact on the health of person. This is especially likely to be greater in the underdeveloped countries where most of the computer users are not aware of the principles and practice of safe computer use 11 Computer vision syndrome along with dry eye syndrome is also one of the major concerns among computer users It is a collection of ocular, 12 extra ocular and systemic symptoms related to prolonged sitting and focusing on computer screens 13 Karachi is a megacity of Pakistan and a large number of people are involved in call centers and software houses. These people work on computers for usually long hours and, thus, are likely to get affected from the computer screens. Literature review has revealed that a few studies have looked into this important computer-related issue 11, 14 therefore, the study aims to assess the frequency of presence of dry eye and its associated risk factors among employees working in call centers and software houses.

Subjects and Methods this cross-sectional study was done during July to October 2014 at Hamdard College of Medicine and Dentistry, Hamdard University, Karachi. The protocol was approved by the Ethical Review Committee of the institute. The data was collected from employees of two call centers and two software houses using convenient sampling technique. All employees of both call center and software houses working on computers were invited to participate in the study. Using the formula for

estimating population proportion with an anticipated prevalence of 67% and 95% confidence interval, 14 the sample size was found to be 156 Six respondents 15 were excluded due to incomplete information hence, a total of 150 employees were included, of both sex and more than 19 years of age. Verbal consent was taken and anonymity of each participant was assured. A pre-designed questionnaire was given to each participant for demographic information and assessment of various eye-related problems. Schirmer's test strips were used to measure production of tears. The strip was placed inside the lower eyelid (conjunctival sac) and the subject was asked to close their eyes for 5 minutes. Both eyes were tested at the same time. The test strip has printed ruler on one side, with a scale divided into millimeters. The strip was removed after 5 minutes and the quantity of tears absorbed was measured and compared with the other eye. To prevent eyes from producing excess tears due to irritation from the paper strips local anesthetic eye drops were given before the start of test. The use of the anesthesia ensures that only basal tear secretion is being measured. The results of Schirmer's test was interpreted as; (i) positive- less than 10 millimeter wetting of the paper strip after 5 minutes, (ii) negative- more than 10 millimeters wetting of the paper strip after 5 minutes. Employees between more than 19 years of age using computer at least for 3 hours daily are included in the study. All participants with any eye morbidities (eye injury, ocular disease), history of ocular surgery, and use of ocular medication or computers less than 3 hours were excluded. The data was entered into SPSS version 20. It was cleaned for data discrepancies and checked for double entries before statistical analysis. Descriptive analysis was done for all categorical variables.

Table 1. Cross-tabulation of demographic characteristics, measures for refractive error correction, type of computer screens used and time spent on computers with presence of eye dryness

Variables	Eye dryness	Total	p-value
Present	n= 61 (%)	n= 89 (%)	
Absent	N= 150 (%)		

Sex

Male	47 (77.0)	70 (78.7)	117 (78.0)
Female	14 (23.0)	19 (21.3)	33 (22.0)

0.843

Age in years

< 25 19 (31.1) 35 (39.3) 54 (36.0)
 > 25 42 (68.9) 54 (60.7) 96 (64.0) 0.387
 Mean ± SD (Range)
 26.9 ± 3.58 (19 to 40)

Wearing spectacles for refractive error correction

Yes 42 (68.9) 35 (39.3) 77 (51.3)
 No 19 (31.1) 54 (60.7) 73 (48.7) <0.0001

Wearing contact lens for refractive error correction

Yes 6 (9.8) 6 (6.7) 12 (8.0)
 No 55 (90.2) 83 (93.3) 138 (92.0) 0.549

Type of computer screen

LCD 52 (85.2) 81 (91.0) 133 (88.7)
 Monitor 9 (14.8) 8 (9.0) 17 (11.3) 0.303

Time spent on computers (in hours)

< 6 26 (42.6) 37(41.6) 63 (42.0)
 > 6 35 (57.4) 52 (58.4) 87 (58.0) 0.999

Table 2.

Association of ocular and extra ocular symptoms with Schirmer's test Variable
 Schirmer's test Total p-value
 Positive Negative N= 150 (%)
 n= 42 (%) n= 108 (%)

Burning of eyes

Yes: 40 (95.2) 26 (24.1) 66 (44.0)
 No: 2 (4.8) 82 (75.9) 84 (56.0) <0.0001

Redness of eyes

Yes: 22 (52.4) 40 (37.0) 62 (41.3)
 No: 20 (47.6) 68 (63.0) 88 (58.7) 0.099

Watery discharge from eyes

Yes: 4 (9.5) 12 (11.1) 16 (10.7)
 No: 38 (90.5) 96 (88.9) 134 (89.3) 0.999

Foreign body sensation in eye

Yes: 1 (2.4) 14 (13.0) 15 (10.0)
 No: 41 (97.6) 94 (87.0) 135 (90.0) 0.068

Dryness of nose

Yes: 17 (40.5) 37 (34.3) 54 (36.0)
 No: 25 (59.5) 71 (65.7) 96 (64.0) 0.570

Dryness of mouth

Yes: 16 (38.1) 36 (33.3) 52 (34.7)
 No: 26 (61.9) 72 (66.7) 98 (65.3) 0.574

Demographic variables were tabulated as frequencies and percentages. Cross-tabulation between different variables and dryness of eyes was done. The frequency of various eye complaints was determined and Chi square test was used to observe the association with Schirmer's test. P-value<0.05 was considered significant.

RESULTS

Out of the total subjects, the majority 117(78%) were males and 33 (22%) were female

participants. The mean age was 26.90 ± 3.58 years (range 19 to 40 years). There were 54 (36%) participants equal to or less than 25 years of age and 96 (64%) participants above 25 years of age. Slightly more than half, 77 (51.3%) of the participants were using spectacles and 12 (8.0%) wore contact lenses for correction of refractive errors. Most of the participants, 87 (58.0%), spent more than 6 hours on computers with the majority, 133(88.7%), having liquid crystal display (LCD) as their computer screens. Statistically significant (p<0.05) association was observed between participants using corrective spectacles with complaint of eye dryness (Table 1).Forty-two (28.0%) participants had positive Schirmer's test (less than 10 millimeters of moisture) while 108 (72%) participants showed normal tearing (more than 10 millimeters of moisture).Burning of eyes, 66 (44%), was the most frequently reported eye complaint among the computer users. This was followed by redness of eye 62 (41.3), watery discharge from eye 16 (10.7%) and 15 (10%) foreign body sensation in eye. Extra ocular symptoms include dryness of nose 54(36%) and dryness of mouth 52 (34.7%) and showed no association with Schirmer's test, whereas burning of eyes was strongly associated (p<0.05) (Table 2).

DISCUSSION

The present study is probably the first study in Karachi conducted on employees of call Centre sand software houses using computers in their job description exclusively. The presence of dry eyewash found to be 28% among participants. A study conducted in Pakistan among medical students reported a much higher frequency (67%) of dry eye syndrome compared to our study Another local 1 4 study conducted on different students found a frequency of 72% However, a recent study on computer 16 users in Lahore reported 30% frequency of dry eyes A study conducted in our neighboring 1 7 country reported frequency of 78.6% for dry eye syndrome Higher frequency of dry eye syndrome 18 was reported from Malaysia 89.9%, and 74% in 1 9 Nigeria 2 0 The mean age of subjects in our study was 27± 3.5 years. Studies from Malaysia and India reported similar mean age of

31 and 30 years respectively. However, Khalaj et al. conducted a study [21, 22] in which the mean age was 15.83 years. The presence of dry eye among participants more than 25 years of age was (68.9%) and more among male participants (77%) in our study suggesting that this prevalence increases with computer exposure time. A study conducted in India reported that the prevalence of dry eye is (25%) among both male and female [24]. The present study found 57.4% of the participants using computers for more than 6 hours a day.

Akinbinu et al. reported that participants who spend 6 to 8 hours average daily on the computer experienced more symptoms of dry eye syndrome from those spending lesser time. Another study also reported of developing DES among those using computer for more than 5 hours per day. Never [21] the less, Reddy et al. reported more than 2 hours continuous use of computer significantly associated with occurrence of DES [19]. In our study, the ocular symptoms experienced by the participants were burning sensation 44%, followed by redness of eyes 41.3%, discharge from eyes 10.7% and foreign body sensation 10%. While among extra ocular symptoms, dryness of nose 36% was the most common symptom. Burning sensation of eye was also reported among 33% and 32% in study conducted in Karachi and India [1, 4] respectively. Another study conducted in India [25] reported relatively less prevalence of burning sensation as it was experienced by 28.9% of the participants. This is in contrast to a study among call center workers reporting a higher prevalence (54.6%) of burning sensation. Logaraj et al. [27] conducted a study in Chennai reported symptoms of redness, burning sensation, headache, and dry eye, and neck and shoulder pain among computer users. In another study, the most common visual [18] problem reported was burning sensation and dry/tired/sore eyes. Mahalingam et al. also reported [28] headache, fatigue and tiredness, burning of eyes, dry eyes, and neck and shoulder pain as the most common symptoms. Prolonged time spent on [29] computer could probably be one of its causes. One of the limitations of this study is that detailed examination of those who had eye complaints was not done due to time restraint and

unavailability of specific machines. More studies should be done among students and professionals involving logistics and experts to identify the burden of this problem that can further avoid complications and limit disability.

CONCLUSION

The present study showed a sizable frequency of dryness of eyes and other ocular and extra ocular symptoms among computer users of call Centre and software houses. There is a definite need to provide them a better understanding of sign and symptoms of dry eye syndrome. Periodic eye examination would help early recognition of computer related eye problems hence, preventing loss of work productivity and vision-related quality of life among its users.

Conflict of Interest

Authors have no conflict of interests and no grant/funding from any organization for this study.

REFERENCES

1. Gayton JL. Etiology, prevalence, and treatment of dry eye disease. *Clin Ophthalmol* 2009;3:405-12.
2. Javadi MA, Feizi S. Dry eye syndrome. *J Ophthalmic Vis Res* 2011;6:192-8.
3. Dilly P. Structure and function of the tear film. In Sullivan DA editor. *Lacrimal Gland, Tear Film, and dry Eye Syndromes*. 1st ed. New York: Springer;1994. p. 239-47.
4. Rolando M, Zierhut M. The ocular surface and tear film and their dysfunction in dry eye disease. *Surv Ophthalmol* 2001;45:S203-10.
5. Sharma A, Hindman HB. Aging: a predisposition to dry eyes. *J Ophthalmol* 2014;2014:781683. [DOI: 10.1155/2014/781683].
6. Schaumberg DA, Uchino M, Christen WG, Semba RD, Buring JE, Li JZ. Patient reported differences in dry eye disease between men and women: impact, management, and patient satisfaction. *PloS One* 2013;8:e76121. [DOI: 10.1371/journal.pone.0076121].
7. Schaumberg DA, Dana R, Buring JE,

- Sullivan DA. Prevalence of dry eye disease among US men: estimates from the Physicians' Health Studies Arch Ophthalmol 2009;127:763-8. [DOI: 10.1001/archophthalmol.2009.103].
8. Chia EM, Mitchell P, Rochtchina E, Lee AJ, Maroun, Wang JJ. Prevalence and associations of dry eye syndrome in an older population: the Blue Mountains Eye Study. Clin Exp Ophthalmol 2003;31:229-32.
 9. Lee AJ, Lee J, Saw SM, Gazzard G, Koh DWidjaja D, et al. Prevalence and risk factors associated with dry eye symptoms: a population based study in Indonesia. Br J Ophthalmol 2002;86:1347-51.
 10. Sahai A, Malik P. Dry eye: prevalence and attributable risk factors in a hospital-based population. Indian J Ophthalmol 2005;53:87-91.
 11. Khan R, Surti A, Rehman R, Ali U. Knowledge and practices of ergonomics in computer users. J Pak Med Assoc 2012;62:213-7.
 12. Uchino M, Schaumberg DA, Dogru M, Uchino Y, Fukagawa K, Shimmura S, et al. Prevalence of dry eye disease among Japanese visual display terminal users. Ophthalmology 2008;115:1982-8. [DOI: 10.1016/j.ophtha.2008.06.022].
 13. Gowrisankaran S, Sheedy JE. Computer vision syndrome: A review. Work 2015;52:303-14. [DOI:10.3233/WOR-152162].
 14. Noreen K, Batool Z, Fatima T, Zamir T. Prevalence of computer vision syndrome and its associated risk factors among undergraduate medical students [Online]. Pak J Ophthalmol 2016;32:1406. Available from: <http://www.pjo.com.pk/32/3/3.%20OA-Khola%20Noreen.pdf>. Accessed on August 21, 2017.
 15. Raosoft sample size calculator [Online]. Raosoft: 2004. Available from: <http://www.raosoft.com/samplesize.html>. Accessed on August 21, 2017
 16. Hassan HMJ, Ehsan S, Arshad HS. Frequency of computer vision syndrome & ergonomic practices among computer engineering students [Online]. International Journal of Science & Research 2016;5:121-5. Available from: <https://www.ijsr.net/archive/v5i5/NOV163207.pdf>. Accessed on August 21, 2017.
 17. Malik AM, Shah SU, Manzoor M. Incidence of dry eyes amongst the computer users. Journal of CMH Lahore Medical College 2017;1:24-27.
 18. Logaraj M, Madhupriya V, Hegde S. Computer vision syndrome and associated factors among medical and engineering students in Chennai Ann Med Health Sci Res 2014;4:179-85. [DOI:10.4103/2141-9248.129028].
 19. Reddy SC, Low CK, Lim YP, Low LL, Mardina F, Nursaleha MP. Computer vision syndrome: a study of knowledge and practices in university students. Nepal J Ophthalmol 2013;5:161-8. [DOI:10.3126/nepjoph.v5i2.8707].
 20. Akinbinu TR, Mashalla YJ. Knowledge of computer vision syndrome among computer users in the workplace in Abuja, Nigeria [Online]. Journal of Physiology and Pathophysiology 2013;4:58-63. Available from: <http://www.academicjournals.org/journal/JPA/article-full-text-pdf/497CABC15892>. Accessed on August 21, 2017. [DOI: 10.5897/JPA.13.0078].
 21. Rahman ZA, Sanip S. Computer user: demographic and computer related factors that predispose user to get computer vision syndrome [Online]. Int J Bus Humanit Technol 2011;1:84-91. Available from: <http://www.ijbhtnet.com/journal/in-dex/43>. Accessed on August 21, 2017.
 22. Arumugam S, Kumar K, Subramani R, Kumar S. Prevalence of Computer Vision Syndrome among Information Technology Professionals Working in Chennai [Online]. World Journal of Medical Sciences 2014;11:312-4. Available from: http://ijbhtnet.com/journals/Vol_1_No_2_September_2011/11.pdf. Accessed on August 21, 2017.
 23. Khalaj M, Ebrahimi M, Shojai P, Bagherzadeh R, Sadeghi T, Ghalenoei M. Computer vision syndrome in eleven to

- eighteen-year-old students in Qazvin [Online]. *Biotech Health Sci* 2015;2:e28234. Available from: https://www.researchgate.net/publication/280533055_Computer_Vision_Syndrome_in_Eleven_to_EighteenYear-Old_Students_in_Qazvin. Accessed on August 21, 2017.
24. Patil SD, Trivedi HR, Parekh NV, Jethva JJ. Evaluation of dry eye in computer users [Online]. *Int J Community Med Public Health* 2016;3:3403-7. Available from: <http://www.ijcmph.com/index.php/ijcmph/article/viewFile/101/99>. Accessed on August 21, 2017.
25. Klamm J, Tarnow KG. Computer Vision Syndrome: A Review of Literature. *Med surg Nurs* 2015;24:89-93.
26. Talwar R, Kapoor R, Puri K, Bansal K, Singh S. A Study of Visual and Musculoskeletal Health Disorders among Computer Professionals in NCR Delhi. *Indian J Community Med* 2009;34:326-8. [DOI: 10.4103/0970-0218.58392].
27. Sa EC, Ferreira Junior M, Rocha LE. Risk factors for computer visual syndrome (CVS) among operators of two call centers in São Paulo, Brazil. *Work* 2012;41:3568-74. [DOI: 10.3233/WOR-20120636-3568].
28. Shantakumari N, Eldeeb R, Sreedharan J, GopalK. Computer use and vision-related problems among university students in Ajman, United Arab Emirate. *Ann Med Health Sci Res* 2014;4:258-63. DOI: 10.4103/2141-9248.129058].
29. Mahalingam. Computer vision syndrome among nurses [Online]. *Journal of Nursing Education and Practice* 2015;1:1-5. Available from: http://www.ramauniversityjournal.com/nursing/pdf_file/1-5.pdf. Accessed on August 21, 2017.