

Research Article

Assessment of hypoalbuminemia in cases of ischemic stroke

**¹Muhammad Javed Masood, ²Trum Noor
and ³Ghulam Mohiuddin**

¹Assistant Professor, Department of Pathology,
Nishter Medical College, Multan

²Assistant Professor, Department of Pathology,
Quaid-e-Azam Medical College, Bahawalpur

³Medical Officer, Basic Health Unit, TalibWala,
Sargodha Cell: 03034064858
Email: drmohiuddin13@gmail.com

ABSTRACT

OBJECTIVES: To assess the hypoalbuminemia in cases of ischemic stroke.

Material and methods: In this cross-sectional study total 250 patients of ischemic stroke both male or female with age range from 40 years to 60 years were recruited from the Department of Medicine, Nishter Medical College, Multan from November 2014 to May 2015.

RESULTS: Mean age of the patients was 48.45 ± 4.675 years. Hypoalbuminemia was noted in 108 (43%) patient. Male patients were 158 (63.2%) and female patients were 92 (36.8%). Total 103 (41.2%) patients belonged to age group 40-50 years and 147 (58.8%) patients belonged to age group 51-60 years, 87 (34.8%) patients were hypertensive, insignificant association of hypoalbuminemia with age, gender and hypertension was seen.

CONCLUSION: Results of this study revealed that higher number of patients found with hypoalbuminemia, male patients were more victim of ischemic stroke as compare to female patient but the difference was not significant. Results of this study also revealed that there is insignificant association of hypoalbuminemia with age and hypertension.

KEY WORDS: Hypoalbuminemia, hypertension, ischemic stroke.

INTRODUCTION:

A stroke is the loss of functions of brain due to disturbance in supply of blood towards the brain. This disturbance of blood supply towards brain may due to hemorrhage or ischemia.¹ Ischemia is caused by either arterial embolism or vessel blockage via thrombosis or by cerebral hypoperfusion.² Almost 800,000 individuals suffer from strokes every year in USA, 82% to 92% of these strokes are ischemic.³ Furthermore, 20% to 40% of cases with ischemic infarction may

develop hemorrhagic transformation within the one week after ictus.⁴

Differentiating between these stroke type is very important because the management of each stroke type is vastly different.⁵ Hypoalbuminemia is a predictive factor for several clinical outcomes (recurrences, functional recovery and medical complications) and mortality in patients with stroke.⁶ Low serum albumin level is frequently

found in hospitalized patients. Hypoalbuminemia was reported in up to 19% of stroke patients.⁷

The exact frequency of Hypoalbuminemia in patients with ischemic stroke is not known as insufficient local data exists.

So the objective of our study is to find out the frequency of hypoalbuminemia in patients with ischemic stroke. Results of this may help us to determine the exact magnitude of this problem which may guide us in better management to decrease the mortality and morbidity related to it.

MATERIAL AND METHODS:

This cross-sectional study was conducted at Department of Medicine, Nishter Medical College, Multan from November 2014 to May 2015. Total 250 patients with ischemic stroke either male or female having age from 40 to 60 years were selected. Patients with decompensated cirrhosis of liver (ultrasound findings of cirrhosis, portal hypertension, ascites, splenomegaly), Nephrotic syndrome (proteinuria (>3.5 gm/day), hypo albuminemia (<3.5g/dl), hypercholesterolemia>200mg/dl and pitting edema.), Hematoma on CT scan brain and Protein losing enteropathy were excluded from the study.

Ischemic stroke was defined as: Patients having hypodense area (infarction) on plain CT scan Brain in the respective vascular territory along with anyone of these: Abnormal reflexes, inability to speak, decreased sensation, loss of balance, mental function problems (irritability and behavioural changes), vision changes (decreased visual acuity <6/6), walking problems (power < 5/5) and weakness within 36 hours of onset.

Study is approved ethically by institutional review board. Written informed consent was taken from every patient.

Five ml fasting blood sample within 36 hours of onset of stroke was drawn and send to the laboratory for serum albumin. Findings were noted in term of hypoalbuminemia (Yes/No) on predesigned proforma along with demographic profile of the patients.

Hypoalbuminemia was labelled when serum albumin level <35 g/l.

Collected data was analyzed by using SPSS version 20.

Mean \pm SD was calculated for age and duration of symptoms as quantitative variable. Qualitative variable like gender, hypertension and hypoalbuminemia was presented as frequencies and percentage. Stratification was done for age, gender, hypertension and duration of symptoms. Post stratification chi-square test was applied to see the effect of these on outcome variable i.e.hypoalbuminemia. P-value \leq 0.05 was considered as significant.

RESULTS:

Total 250 patients with ischemic stroke were selected to evaluate the hypoalbuminemia. Mean age of the patients was 48.45 ± 4.675 years. Among the 250 patients hypoalbuminemia was found in 108 (43%) patients. (Fig. 1)

Patients were divided into two age groups, age group 40-50 years and age group 51-60 years. There were 103 (41.2%) patients in age group 40-50 years and 147 (58.8%) patients in age group 51-60 years and hypoalbuminemia was found in 48 (46.6%) and 60 (40.8%) patients respectively.

Insignificant (P = 0.367) association between age and hypoalbuminemia was noted. (Table 1).

Out of 158 (63.2%) male patients, hypoalbuminemia was noted in 70 (44.3%) patients and out of 92 (36.8%) female patients, hypoalbuminemia was noted in 38 (41.3%) patients. Insignificant (P = 0.692) association of hypoalbuminemia with gender was noted. (Table 2)

As shown in table 3, out of 87 (34.8%) hypertensive patients, hypoalbuminemia was seen in 42 (48.28%) patients and out of 163 (65.5%) normotensive patients, hypoalbuminemia was seen in 66 (40.5%) patients. Insignificant (P = 0.283) association was seen between hypertension and hypoalbuminemia was noted.

Fig. 1: Frequency for Hypoalbuminemia




Table 1: Stratification for age

Age	Hypoalbuminemia		Total	P. value
	Yes (%)	No (%)		
40-50	48 (46.6)	55 (53.4)	103 (41.2)	0.367
51-60	60 (40.8)	87 (59.2)	147 (58.8)	
Total	108 (43)	142 (57)	250	

Table 2: Stratification for Gender

Gender	Hypoalbuminemia		Total	P. value
	Yes (%)	No (%)		
Male	70 (44.3)	88 (55.7)	158 (63.2)	0.692
Female	38 (41.3)	54 (58.7)	92 (36.8)	
Total	108 (43)	142 (57)	250	

Table 3: Stratification for hypertension

Hypertension	Hypoalbuminemia		Total	P. value
	Yes (%)	No (%)		
Yes	42 (48.28)	45 (51.72)	87 (34.8)	0.283
No	66 (40.5)	97 (59.5)	163 (65.5)	
Total	108 (43)	142 (57)	250	

DISCUSSION:

In west, stroke affect the 700000 people every year and in hospital mortality rate is 3-5% according to type of stroke.^{8,9} Generally it is believed that early death after stroke is mainly attributable to the disease itself, whereas the death after acute phase is due to the hospitalization and the related complications during this period.^{10,11}

This study showed frequency of hypoalbuminemia as 43% in patients of ischemic stroke. Similarly Dziedzic et al⁷ reported frequency of hypoalbuminemia as 45.5% patients. Vahedi A et al.,¹² reported frequency of hypoalbuminemia in 43% patients of ischemic stroke. Chen Y et al.,¹³ studied serum albumin levels in 70 patients of ischemic stroke and observed hypoalbuminemia in 56% patients.

In our study, we found a higher number of patients with hypoalbuminemia. Some other studies reported lower frequency of hypoalbuminemia in patients of ischemic stroke. In one study Davalos et al.¹⁴ found hypoalbuminemia only in 7.7% patients out of 104 patients of ischemic stroke and Gariballa et al.,¹⁵ reported hypoalbuminemia in 19% patients. Davis et al.,¹⁶ found hypoalbuminemia in 16.2% patients and Dzieniszewski et al.,¹⁷ hypoalbuminemia in 20.7% patients.

CONCLUSION:

Results of this study revealed that higher number of patients found with hypoalbuminemia, male patients were more victim of ischemic stroke as compare to female patient but the difference was not significant. Results of this study also revealed that there is insignificant association of hypoalbuminemia with age and hypertension.

REFERENCES:

1. Sims NR, Muyderman H. Mitochondria, oxidative metabolism and cell death in stroke. *Biochimica et Biophysica Acta*. 2009;1802(1):80–91.
2. Fonarow GC, Saver JL, Smith EE, Broderick JP, Kleindorfer DO, Sacco RL, et al.

- Relationship of national institutes of health stroke scale to 30-day mortality in medicare beneficiaries with acute ischemic stroke. *J Am Heart Assoc.* 2012;1(1):42-50.
3. Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, et al. Heart disease and stroke statistics--2012 update: a report from the American Heart Association. *Circulation.* 2012;125(1):e2-e220.
 4. Mullins ME, Lev MH, Schellingerhout D, Gonzalez RG, Schaefer PW. Intracranial hemorrhage complicating acute stroke: how common is hemorrhagic stroke on initial head CT scan and how often is initial clinical diagnosis of acute stroke eventually confirmed?. *AJNR Am J Neuroradiol.* Oct 2005;26(9):2207-12.
 5. Nighoghossian N, Hermier M, Adeleine P, Blanc-Lasserre K, Derex L, Honnorat J. Old microbleeds are a potential risk factor for cerebral bleeding after ischemic stroke: a gradient-echo T2-weighted brain MRI study. *Stroke.* Mar 2002;33(3):735-42.
 6. Lázaro VA, del SerQuijano T, Martín RB. Hypoalbuminemia and other prognostic factors of mortality at different time points after ischemic stroke. *Nutr Hosp.* 2013;28(2):456-63.
 7. Dziedzic T, Pera J, Slowik A, Gryz-Kurek EA, Szczudlik A. Hypoalbuminemia in acute ischemic stroke patients: frequency and correlates. *Eur J Clin Nutr.* 2007;61(11):1318-22.
 8. Members WG, Thom T, Haase N, Rosamond W, Howard VJ, Rumsfeld J, et al. Heart Disease and Stroke Statistics—2006 Update A Report From the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation.* 2006 Feb 14;113(6):e85-151.
 9. Liebeskind DS, Kidwell CS, Sayre JW, Saver JL. Evidence of publication bias in reporting acute stroke clinical trials. *Neurology.* 2006 Sep 26;67(6):973-9.
 10. Famakin B, Weiss P, Hertzberg V, McClellan W, Presley R, Krompf K, et al. Hypoalbuminemia predicts acute stroke mortality: Paul Coverdell Georgia Stroke Registry. *J Stroke Cerebrovasc Dis.* 2010 Jan;19(1):17-22.
 11. Ebersole JL, Cappelli D. Acute-phase reactants in infections and inflammatory diseases. *Periodontol 2000.* 2000 Jun;23:19-49.
 12. Vahedi A, Lotfinia I, Sad RB, Halimi M, Baybordi H. Relationship between admission hypoalbuminemia and inhospital mortality in acute stroke. *Pak J Biol Sci.* 2011 Jan 15;14(2):118-22.
 13. Chen Y, Li S, Sun F, Liu Y, Hu W. Monitoring of medical complications after acute ischemic stroke in a neurological intensive care unit. *Eur Neurol.* 2011;66(4):204-9.
 14. Dávalos A, Ricart W, Gonzalez-Huix F, Soler S, Marrugat J, Molins A, et al. Effect of Malnutrition After Acute Stroke on Clinical Outcome. *Stroke.* 1996 Jun 1;27(6):1028-32.
 15. Gariballa SE, Parker SG, Taub N, Castleden CM. Influence of nutritional status on clinical outcome after acute stroke. *Am J Clin Nutr.* 1998 Aug;68(2):275-81.
 16. Davis JP, Wong AA, Schluter PJ, Henderson RD, O'Sullivan JD, Read SJ. Impact of premorbid undernutrition on outcome in stroke patients. *Stroke.* 2004 Aug;35(8):1930-4.
 17. Dzieniszewski J, Jarosz M, Szczygieł B, Długosz J, Marlicz K, Linke K, et al. Nutritional status of patients hospitalised in Poland. *Eur J Clin Nutr.* 2005 Apr;59(4):552-60.