

Research Article

Phytotherapy for *Streptococcus sanguis*

**Somayeh Delfani^{1,2}, Faranak Rezaei^{1,2*}
and Somayeh Shahsavari³**

¹Department of Microbiology, School of Medicine,
Lorestan University of Medical Sciences, Khorramabad, Iran

²Razi Herbal Medicines Research Center,
Lorestan University of Medical Sciences, Khorramabad, Iran

³Biotechnology and Medicinal Plants Research Center,
Ilam University of Medical Sciences, Ilam, Iran

Corresponding author. Dr. Faranak Rezaei; Department of Microbiology,
School of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

ABSTRACT

Streptococcus mutans and *Streptococcus sanguis* are the most important bacterial agents of dental caries. *Streptococcus sanguis* is the most commonly isolated species of the dental plaque. Medicinal plants are rich sources of bioactive and antioxidant substances which treat various diseases with various mechanisms. Since *Streptococcus sanguis* is one of the bacterial agents responsible for dental caries, the efforts in this article were on reporting the medicinal plants which have an effective effect on this bacteria. Searching for articles with words of *Streptococcus sanguis*, phytotherapy, medicinal herbs, and tooth diseases was done in present reviewing study. Searching was done from databases such as Scopus, ISC, SID, Meiran, and some other databases.

Accordingly, 23 drugs which are effective on *Streptococcus sanguis* are used. It was determined based on the obtained results that *Cinnamomum cassia*, *Stachys byzanthina*, *Stachys lavandulifolia*, *Phlomis brugueur*, *Juglans regia*, *Lonicera japonica*, *Jasminum sambac*, *Eucommia ulmoides*, *Camellia sinensis*, *Gynostemma pentaphyllum*, *Cymbopogon citratus*, *Ilex paraguariensis*, *Rosmarinus officinalis*, *Kaempferia pandurata*, *Azadirachta indica*, *Acacia nilotica*, *Ocimum basilicum*, *Emblica officinalis*, *Terminalia bellirica*, *Syzygium cumini*, *Syzygium aromaticum*, *Diclinanona calycina* and *Thyrsodium spruceanum* medicinal plants are the most important medicinal plants with anti-*Streptococcus sanguis* properties. Medicinal plants used in this study have anti-*Streptococcus sanguis* properties due to presents of bioactive substances, antioxidants, flavonoids, flavones, anthocyanins, and the like.

Keywords: Medicinal plants, *Streptococcus sanguis*

INTRODUCTION

Streptococci are the most important gram-positive bacteria that can cause toxic septicemia and childhood infections. These bacteria are genetically diverse which contain important human pathogens (1).

Some bacteria are the main cause of dental caries (2,3). *Streptococcus mutans* and *Streptococcus sanguis* are the most important bacterial agents of dental caries (4). *Streptococcus sanguis* is the most commonly isolated species of the dental plaque that it is the major causative factor of periodontitis (5).

This bacteria provides the energy necessary to grow the bacteria in the absence of carbohydrate fermentation via the mechanism of hydrolysis of arginine. Hence, bacteria can survive and function without the existence of carbohydrate fermentation (4).

The efforts have been on stopping the growth of bacteria causing dental decay by using plaque control methods such as toothbrushes and the use of antibacterial mouthwashes but if all of these arrangements do not work, the accumulation of these microorganisms will destroy the ivory and enamel and will eventually lead to loss of teeth (6, 7).

Medicinal plants have been used to treat dental diseases since ancient times (8-11). Medicinal plants are rich sources of bioactive and antioxidant substances (12-14) which will treat various diseases with various mechanisms (15-18).

Since *Streptococcus sanguis* is one of the bacterial agents responsible for dental caries, the efforts in this study were on reporting medicinal plants that have an effective effect on this bacteria.

METHOD OF STUDY

Searching for articles with words of *Streptococcus sanguis*, phytotherapy, medicinal herbs, and tooth diseases was done in present reviewing study. Searching was done from databases such as Scopus, ISC, SID, Magiran, and some other databases.

RESULTS

It was determined based on the obtained results that *Cinnamomum cassia*, *Stachys byzanthina*, *Stachys lavandulifolia*, *Phlomis brugueur*, *Juglans regia*, *Lonicera japonica*, *Jasminum sambac*, *Eucommia ulmoides*, *Camellia sinensis*, *Gynostemma pentaphyllum*, *Cymbopogon citratus*, *Ilex paraguariensis*, *Rosmarinus officinalis*, *Kaempferia pandurata*, *Azadirachta indica*, *Acacia nilotica*, *Ocimum basilicum*, *Emblica officinalis*, *Terminalia bellirica*, *Syzygium cumini*, *Syzygium aromaticum*, *Diclinanone calycina* and *Thyrsodium spruceanum* medicinal plants are the most important medicinal plants with anti-*Streptococcus sanguis* properties. Further details in this regard are provided in Table 1.

Phytotherapy for *Streptococcus sanguis*

No	Scientific Name	Part of plant	Family Name	Common Name	Origin of plant	Country Of study	year	Bacteria	Result	ref
1	CINNAMOMUM CASSIA	Bark	Lauraceae	Chinese cinnamon	Pakistan	Pakistan	In vitro 2006	human isolates of <i>Streptococcus sanguis</i>	The diameter of inhibition zone of oil of this plant for bacteria was reported to be 27 mm	(19)
2	Stachysbyzanthina		Lamiaceae	lamb's-ear	iran	iran	In vitro 1386	<i>Streptococcus sanguis</i>	Antibacterial properties of methanolic extract of this plant were proven with a concentration of 1000 µg / ml through the diameter of the inhibition zone of 15 mm. However, the diameter of inhibition zone for gentamicin and amikacin halo was respectively 24 mm and 19 mm	(20)
3	Stachyslavandulifolia		Lamiaceae		iran	iran	In vitro 1386	<i>Streptococcus sanguis</i>	Antibacterial properties of methanolic extract of this plant were proven with a concentration of 1000 µg / ml through the diameter of the inhibition zone of 15 mm. However, the diameter of inhibition zone for gentamicin and amikacin halo was respectively 24 mm and 19 mm	(20)
4	Phlomisbruguieri		Lamiaceae		iran	iran	In vitro 1386	<i>Streptococcus sanguis</i>	Antibacterial properties of methanolic extract of this plant were proven with a concentration of 1000 µg / ml through the diameter of the inhibition zone of 18 mm. However, the diameter of inhibition zone for gentamicin and amikacin halo was respectively 24 mm and 19 mm	(20)
5	Juglansregia	Leaf	Juglandaceae	English walnut	iran	iran	In vitro 1389	<i>Streptococcus sanguis</i> (PTCC:1449)	Antibacterial properties of methanolic extract of walnut leaves was proven with MIC = 15.6µg / ml and MBC = 31.25µg / ml. while, chlorhexidine as an antibiotic control had MIC = 15.6 and MBC = 31.25.	(21)
6	Lonicera japonica	Flowers	Caprifoliaceae	Honeysuckle	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 4 mg / ml inhibited the growth of this bacteria	(22)
7	Jasminumsambac	Flowers	Oleaceae	Jasmine	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 1 mg / ml inhibited the growth of this bacteria	(22)
8	Eucommiaulmoides	Leaves	Eucommiaceae	Duzhong	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 4 mg / ml inhibited the growth of this bacteria	(22)
9	Camellia sinensis	Leaves	Theaceae	Green tea	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 4 mg / ml inhibited the growth of this bacteria	(22)
10	Gynostemma pentaphyllum	Leaves	Cucurbitaceae	Jiaogulan	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 1 mg / ml inhibited the growth of this bacteria	(22)
11	Cymbopogon citratus	Leaves	Poaceae	Lemon grass	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 1 mg / ml inhibited the growth of this bacteria	(22)

Phytotherapy for *Streptococcus sanguis*

12	<i>Ilex paraguariensis</i>	Leaves	Aquifoliaceae	Yerba mate	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 4 mg / ml inhibited the growth of this bacteria	(22)
13	<i>Rosmarinus officinalis</i>	Leaves	Lamiaceae	rosemary	Taiwan	Taiwan	In vitro 2008	<i>Streptococcus sanguinis</i> (BCRC15273)	The methanolic extract of this plant with MIC = 2 mg / ml inhibited the growth of this bacteria	(22)
14	<i>Kaempferiapandurata</i>	Rhizome	Zingiberaceae		Indonesia	South Korea	In vitro 2004	<i>Streptococcus sanguis</i> ATCC 35105	The antibacterial properties of isopanduratin A composition existing in this plant was proven with MIC = 4 mg / ml and MBC = 8 mg / ml.	(23)
15	<i>Azadirachta indica</i>	Foliage	Meliaceae	Neem	India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 10.6 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 0 mm for nitrofurantoin	(24)
16	<i>Acacia nilotica</i>	Foliage	Fabaceae	gum arabic tree	India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 11.3 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 11.3 mm for carbenicillin, 0 mm for nitrofurantoin and 11.3 mm for tetracycline	(24)
17	<i>Ocimum basilicum</i>	Foliage	Lamiaceae	Basil	India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 10.6 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 0 mm for nitrofurantoin	(24)
18	<i>Emblica officinalis</i>	Foliage	Euphorbiaceae		India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 29.6 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 12 mm for gentamicin , 11.3 mm for carbenicillin, 0 mm for nitrofurantoin and 11.3 mm for tetracycline	(24)
19	<i>Terminalia bellirica</i>	Foliage	Combretaceae	Beleric	India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 28.6 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 12 mm for gentamicin , 11.3 mm for carbenicillin, 12.6 mm for nalidixic acid, 0 mm for nitrofurantoin and 11.3 mm for tetracycline	(24)

Phytotherapy for *Streptococcus sanguis*

20	<i>Syzygiumcumini</i>	Foliage	Myrtaceae	jambul	India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 25 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 12 mm for gentamicin , 11.3 mm for carbenicillin, 12.6 mm for nalidixic acid, 0 mm for nitrofurantoin and 11.3 mm for tetracycline	(24)
21	<i>Syzygiumaromaticum</i>	Floral parts	Myrtaceae	Cloves	India	India	In vitro 2012	Clinical isolates of <i>Streptococcus sanguis</i>	Antibacterial properties of the aqueous extract of this plant was proven by having a diameter of 23.6 mm inhibition zone. However, the diameter of the inhibition zone for antibiotic controllers of ampicillin was 10.6 mm and it was 12 mm for gentamicin , 11.3 mm for carbenicillin, 12.6 mm for nalidixic acid, 0 mm for nitrofurantoin and 11.3 mm for tetracycline	(24)
22	<i>Diclinanonalcalycina</i>	Leaf	Annonaceae		Brazil	Brazil	In vitro 2008	<i>Streptococcus sanguis</i> (ATCC 15300)	Antibacterial property of the chloropharmaceutical extract of this plant was with having a diameter of 22 mm for inhibition zone. MIC of this plant was also 48.8 µg / ml. while the diameter of the inhibition zone of rifampin as control was 20 mm	(25)
23	<i>Thyrsodiumspruceanum</i>	Branch	Anacardiaceae		Brazil	Brazil	In vitro 2008	<i>Streptococcus sanguis</i> (ATCC 15300)	Antibacterial property of the methanolic extract of this plant was with having a diameter of 20 mm for inhibition zone. MIC of this plant was also 48.8 µg / ml. while the diameter of the inhibition zone of rifampin as control was 20 mm	(25)

Table 1. Medicinal plants affecting *Streptococcus sanguis*

DISCUSSION

Medicinal plants used in this study have anti-*Streptococcus sanguis* properties due to presents of bioactive substances, antioxidants, flavonoids, flavones, anthocyanins, and the like. Although the mechanism actions of these plants are not clear, but with no debt, these by-products are responsible for their actions. From these, phenolic components of these plants in various investigations have been shown to possess antimicrobial activity, other than having other benefits (26-35). The plants presented in this review and a lot of other plants have phenolic compounds (37-40). These plants, therefore, which possess phenolic compounds, might also be effective against pathogenic microorganism.

The resistance of microbial infections to anti-biotic agents has been increasing in the recent years Multi-drug resistance has huge challenges and infections due to multi-resistant bacteria especially in the intensive care units impose more serious problems. Medicinal plants also have been shown to be effective on resistant bacteria. More importantly, the microorganism mechanism actions and resistance are different (41-52). Medicinal plants have various active compounds which each one may have its own effect. Therefore, they may act by various mechanisms to overcome on microorganisms or other diseases (53-61). In other words, consumption of medicinal plants to treat or prevent human diseases has certain advantages. They are mostly effective in multiple diseases (62-74). They also are mostly cheap, readily available as well as being less toxic environmental hazards and biodegradable. Effective plant extracts are able to combat human pathogenic microorganisms other than treating other complications which might be associated with infection (75-84). There is therefore, renewed interest for medicinal plants to search for plants with anti-microbial activities, especially the plants which have the potential to treat other diseases (85-88). This has led to discovery of plants with low side effects and high efficacy.

REFERENCES

1. Hon KL, Fu A, Leung TF, Poon TC, Cheung W CY, et al. Cardiopulmonary morbidity of streptococcal infections in a PICU. The clinical respiratory journal. 2015;9(1):45-52.
2. Maghareabed A, Bateni E, Rabiei A, Pour Moradi B. The Review on the Effect of Using Mouthwash in Keeping Mouth and Teeth Healthy. Isfahan Dentistry Faculty journal. 2012;7(5).
3. Ambrosio SR, Furtado NA, de Oliveira DC, da Costa FB, Martins CH, de Carvalho TC. Antimicrobial activity of kauranedi terpenes against oral pathogens. Z Naturforsch C. 2008;63(5-6):326-30.
4. Kuhnert WL, Quivey Jr RG Jr. Genetic and Biochemical Characterization of the F-TPase operon from *Streptococcus sanguis* 10904. J Bacteriol. 2003;185(5):1525-33.
5. Faghri J., Moghim SH., Moghareh Abed A., Rezaei F., Chalabi M. Prevalence of *Porphyromonas gingivalis* and *Bacteroides forsythous* in chronic Periodontitis by multiplex PCR. Pakistan Journal of biological sciences. 2007; 10(22): 4123-4127
6. Lemme A, Gröbe L, Reck M, Tomasch J, Wagner-Döbler I. Sub population Specific Transcriptome analysis of competence - Stimulating- Peptide- Induced *Streptococcus mutans*. J Bacteriol 2011; 193(8):1863-77
7. Yamaguchi M, Terao Y, Ogawa T, Takahashi T, Hamada S, Kawabata S. Role of *Streptococcus sanguinis* sortase A in bacterial colonization. Microbes Infect 2006;8(12-13):2791-6
8. Ghamari S, Mohammadrezaei-Khorramabadi R, Mardani M, Shahsavari S. An Overview of the most Important Medicinal Plants with Anti-Toothache Property based on Ethnobotanical Sources in Iran. J. Pharm. Sci. & Res 2017; 9(6): 796-799.

9. Delfani S, Mohammadrezaei-Khorramabadi R, Abbaszadeh S, Naghdi N, Shahsavari S. Phytotherapy for *Streptococcus pyogenes*. *J. Pharm. Sci. & Res* 2017; 9(5): 513-526.
10. Ghamari S, Abbaszadeh S, Mardani M, Shahsavari S. Identifying medicinal plants affecting the teeth from the Southern District of Ilam province, Iran. *J. Pharm. Sci. & Res* 2017; 9(6): 800-803.
11. Ghamari S, Mohammadrezaei-Khorramabadi R, Mardani M, Shahsavari S. An Overview of the most important Medicinal Plants used as Mouth Freshener. *J. Pharm. Sci. & Res* 2017; 9(6): 804-807.
12. Zarei L, Naji-Haddadi S, Pourjabali M, Naghdi N, Tasbih-Forosh M, Shahsavari S. Systematic Review of Anti-Rheumatic Medicinal Plants: An Overview of the Effectiveness of Articular Tissues and Joint Pain Associated with Rheumatoid Arthritis. *J. Pharm. Sci. & Res. Vol. 9(5)*, 2017, 547-551.
13. Pourjabali M, Mohammadrezaei-Khorramabadi R, Abbaszadeh S, Naghdi N, Naji-Haddadi S, Bahmani F. Medicinal Plants Used For Hypertension. *J. Pharm. Sci. & Res. Vol. 9(5)*, 2017, 537-541.
14. Karimi M, Naghdi N, Naji-Haddadi S, Bahmani F. Medicinal Plants Used For Kidney Pain. *J. Pharm. Sci. & Res. Vol. 9(5)*, 2017, 542-546.
15. Delfani S, Mohammadrezaei-Khorramabadi R, Ghamari S, Khadivi-Boroujeni R, Khodabandelo N, GhadiraliKhorzoughi M, Shahsavari S. Systematic Review for Phytotherapy in *Streptococcus Mutans*. *J. Pharm. Sci. & Res. Vol. 9(5)*, 2017, 552-561.
16. Sadighara P, Erfanmanesh A, Haghi E, Nikaein D, Mohajerfar T, Tohidimoghadam T, Bahmani M, Abbaszadeh A, Rafieian-Kopaei M. The Antibacterial Activity of Chitosan between Different Extraction Method. *Journal of Chemical and Pharmaceutical Science* 2016; 9(3): 1325-1328.
17. Zarei L, Pourjabali M, Naghdi N, Naji-Haddadi S, Bahmani E. A Systematic Review of the Most Important Medicinal Plants Native to Iran Effective on Testicular Morphology and Hormonal Testicular Function. *J. Pharm. Sci. & Res. Vol. 9(5)*, 2017, 562-567.
18. Asadi N, Bahmani M, Shahsavari S and Asadi-Samani M. Identification and Introduction of The Medicinal Plants Used by Honeybees in Markazi Province. *International Journal of Pharmaceutical and Phytopharmacological Research* 2017; 7(2): 15-18.
19. Chaudhry NMA, Tariq P. Anti-microbial activity of *Cinnamomum cassia* against diverse microbial flora with its nutritional and medicinal impacts. *Pakistan Journal of Botany*. 2006;38(1):169.
20. Semnani M., Katayoun S, Mahdavi, Rahimi M. Antimicrobial effect of methanolic extracts of several plant species from *Stachys* and *Phlomis* species. *Journal of Mazandaran University of Medical Sciences*. 2007; 17 (57): 57-66.
21. Chaloshitori SH, rafieian-Kopaei M, Ashrafi D. Comparison of antibacterial effects of ethanolic extract of Iranian walnut leaves with chlorhexidine mouthwash on *Streptococcus mutans* and *Streptococcus sanguinis*. *Dental Journal of Islamic Society of Dentists*. 2010; 22 (4): 211-7.
22. Tsai T-H, Tsai T-H, Chien Y-C, Lee C-W, Tsai P-J. In vitro antimicrobial activities against cariogenic streptococci and their antioxidant capacities: A comparative study of green tea versus different herbs. *Food Chemistry*. 2008;110(4):859-64.
23. Hwang J-K, Chung J-Y, Baek N-I, Park J-H. Isopanduratin A from *Kaempferia pandurata* as an active antibacterial agent against cariogenic *Streptococcus mutans*.

- International journal of antimicrobial agents. 2004;23(4):377-81.
24. Pathak A, Sardar A, Kadam V, Rekadwad B, Karuppayil SM. Efficacy of some medicinal plants against human dental pathogens. *Indian Journal of Natural Products and Resources*. 2012;3(1):123-7.
 25. Carneiro ALB, Teixeira MFS, Oliveira VMAd, Fernandes OCC, Cauper GSdB, Pohlit AM. Screening of Amazonian plants from the Adolpho Ducke forest reserve, Manaus, state of Amazonas, Brazil, for antimicrobial activity. *Memórias do Instituto Oswaldo Cruz*. 2008;103(1):31-8.
 26. Karimi A, Majlesi M, Rafieian-Kopaei M. Herbal versus synthetic drugs; beliefs and facts. *J Nephrofarmacol* 2015; 4(1): 27-30.
 27. Sedighi M, Rafieian-kopaei M, Noori-Ahmadabadi M. Effect of *Allium ampeloprasum* on ileum function: Involvement of beta-adrenergic receptors and voltage dependent calcium channels. *Life Sci J*. 2012;9(4):1660-7. PubMed PMID: WOS:000316682500252.
 28. Rafieian-Kopaei M, Setorki M, Heidarian E, Shahinfard N, Ansari R. Effect of *Anethum graveolens* on hypelipidemia induced hepatotoxicity. *Toxicol Lett*. 2012 Jun 17;211:S167-S. PubMed PMID: WOS:000305173900545.
 29. Hajian S. Positive effect of antioxidants on immune system. *Immunopathol Persa*. 2015;1(1):e02.
 30. Gholamian-Dehkordi N, Luther T, Asadi-Samani M, Mahmoudian-Sani MR. An overview on natural antioxidants for oxidative stress reduction in cancers; a systematic review. *Immunopathol Persa*. 2017;3(2):e12.
 31. Seddighi S, Bahmani M, Asgary S, Beyranvand F, Rafieian-Kopaei M. A review of plant-based compounds and medicinal plants effective on atherosclerosis. *J Res Med Sci*. 2017; 22:30,
 32. Jalaly L, Sharifi G, Faramarzi M, Nematollahi A, Rafieian-Kopaei M, Amiri M, et al. Comparison of the effects of *Crataegus oxyacantha* extract, aerobic exercise and their combination on the serum levels of ICAM-1 and E-Selectin in patients with stable angina pectoris. *Daru*. 2015 Dec 19;23.
 33. Shirzad H, Taji F, Rafieian-Kopaei M. Correlation Between Antioxidant Activity of Garlic Extracts and WEHI-164 Fibrosarcoma Tumor Growth in BALB/c Mice. *J Med Food*. 2011 Sep;14(9):969-74.
 34. Akhlaghi M, Shabanian G, Rafieian-Kopaei M, Parvin N, Saadat M, Akhlaghi M. Citrus aurantium Blossom and Preoperative Anxiety. *Rev Bras Anesthesiol*. 2011 Nov-Dec;61(6):702-12.
 35. Nasri H, Rafieian-Kopaei M. Comment on: The effects of green tea consumption on metabolic and anthropometric indices in patients with type 2 diabetes. *J Res Med Sci*. 2014 Dec;19(12):1203-4.
 36. Sedighi M, Rafieian-kopaei M, Noori-Ahmadabadi M. Efect of *Allium ampeloprasum* on ileum function: Involopersum of beta-adrenergic receptors and voltage depend calcium channels. *Life Science jornal* 2012;9(3):1660-7
 37. Ghaed F, Rafieian-Kopaei M, NematbakhshM, Baradaran A. Ameliorative effects of metformin on renal histologic and biochemical alterations of gentamicin-induced renal toxicity in Wistar rats. *J Res Med Sci*. 2012; 17: 621-5.
 38. Rafieian-Kopaei M, Baradaran A, Merrikhi A, Nematbakhsh M, Madihi Y, Nasri H. Efficacy of co-administration of garlic extract and metformin for prevention of gentamicin renal toxicity in wistar rats: A biochemical study. *Int J Prev Med*. 2013; 4: 258-64.
 39. Shirzad H, Shahrani M, Rafieian-Kopaei M. Comparison of morphine and tramadol effects on phagocytic activity of mice

- peritoneal phagocytes in vivo. *IntImmunopharmacol*. 2009 Jul;9(7-8):968-70.
40. Bagheri N, Azadegan-Dehkordi F, Shirzad M, Zamanzad B, Rahimian G, Taghikhani A, et al. Mucosal interleukin-21 mRNA expression level is high in patients with *Helicobacter pylori* and is associated with the severity of gastritis. *Cent Eur J Immunol*. 2015;40(1):61-7.
 41. Bahmani M, Rafieian-Kopaei M, Parsaei P, Mohsenzadegan A. The anti-leech effect of *Peganumharmala* L. extract and some anti-parasite drugs on *Limnatisnilotica*. *Afr J Microbiol Res*. 2012; 16;6(10):2586-90.
 42. Bagheri N, Taghikhani A, Rahimian G, Salimzadeh L, Dehkordi FA, Zandi F, et al. Association between virulence factors of *Helicobacter pylori* and gastric mucosal interleukin-18 mRNA expression in dyspeptic patients. *Microb Pathogenesis*. 2013;65:7-13.
 43. Shirzad H, Burton RC, Smart YC, Rafieian-Kopaei M, Shirzad M. Natural Cytotoxicity of NC-2(+) Cells Against the Growth and Metastasis of WEHI-164 Fibrosarcoma. *Scand J Immunol*. 2011;73(2):85-90.
 44. Bagheri N, Rahimian G, Salimzadeh L, Azadegan F, Rafieian-Kopaei M, Taghikhani A, et al. Association of the Virulence Factors of *Helicobacter pylori* and Gastric Mucosal Interleukin-17/23 Mrna Expression in Dyspeptic Patients. *Excli J*. 2013;12:5-14.
 45. Bagheri N, Azadegan-Dehkordi F, Sanei H, Taghikhani A, Rahimian G, Salimzadeh L, et al. Associations of a TLR4 single-nucleotide polymorphism with *H. pylori* associated gastric diseases in Iranian patients. *Clin Res Hepatol Gas*. 2014 Jun;38(3):366-71.
 46. Bagheri N, Azadegan-Dehkordi F, Shirzad H, Rafieian-Kopaei M, Rahimian G, Razavi A. The biological functions of IL-17 in different clinical expressions of *Helicobacter pylori*-infection. *Microb Pathogenesis*. 2015 Apr;81:33-8.
 47. Razavi A, Bagheri N, Azadegan-Dehkordi F, Shirzad M, Rahimian G, Rafieian-Kopaei M, et al. Comparative Immune Response in Children and Adults with *H. pylori* Infection. *J Immunol Res*. 2015.
 48. Bahmani M, Banihabib E, Rafieian-Kopaei M, Gholami-Ahangaran M. Comparison of Disinfection Activities of Nicotine with Copper Sulphate in water Containing *Limnatisnilotica*. *KafkasUniv Vet Fak*. 2015 Jan-Feb;21(1):9-11.
 49. Azadegan-Dehkordi F, Bagheri N, Shirzad M, Sanei MH, Hashemzadeh-Chaleshtori M, Rafieian-Kopaei M, et al. Correlation Between Mucosal IL-6 mRNA Expression Level and Virulence Factors of *Helicobacter pylori* in Iranian Adult Patients With Chronic Gastritis. *Jundishapur J Microb*. 2015 Aug;8(8).
 50. Rafieian-Kopaei M, Nasri H, Alizadeh F, Ataei B, Baradaran A. Immunoglobulin A Nephropathy and Malaria *falciparum* Infection; a Rare Association. *Iran J Public Health*. 2013 May;42(5):529-33.
 51. Bahmani M, Saki K, Rafieian-Kopaei M, Karamati SA, Eftekhari Z, Jelodari M. The most common herbal medicines affecting *sarcomastigophora* branches: a review study. *Asian Pac J Trop Med*. 2014 Sep;7:S14-S21.
 52. Asgari S, Setorki M, Rafieian-Kopaei M, Heidarian E, Shahinfard N, Ansari R, et al. Postprandial hypolipidemic and hypoglycemic effects of *Allium hertifolium* and *Sesamumindicum* on hypercholesterolemic rabbits. *Afr J Pharm Pharmacol*. 2012 Apr;6(15):1131-5.
 53. Baradaran A, Nasri H, Rafieian-Kopaei M. Protection of Renal Tubular Cells by Antioxidants: Current Knowledge and New Trends. *Cell J*. 2015;16(4):568-71.
 54. Heidarian E, Rafieian-Kopaei M. Protective effect of artichoke (*Cynarascolymus*) leaf

- extract against lead toxicity in rat. *Pharm Biol.* 2013 Sep;51(9):1104-9.
55. Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci.* 2014 Jan;19(1):82-3.
 56. Baradaran A, Nasri H, Rafieian-Kopaei M. Oxidative stress and hypertension: Possibility of hypertension therapy with antioxidants. *J Res Med Sci.* 2014 Apr;19(4):358-67.
 57. Rafieian-Kopaei M, Baradaran A, Rafieian M. Oxidative stress and the paradoxical effects of antioxidants. *J Res Med Sci.* 2013 Jul;18(7):628-.
 58. Baradaran A, Madihi Y, Merrikhi A, Rafieian-Kopaei M, Nematbakhsh M, Asgari A, et al. Nephrotoxicity of hydroalcoholic extract of *Teucrium polium* in Wistar rats. *Pak J Med Sci.* 2013;29:329-33.
 59. Bahmani M, Zargarani A, Rafieian-Kopaei M, Saki K. Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran. *Asian Pac J Trop Med.* 2014 Sep;7:S348-S54.
 60. Nasri H, Nematbakhsh M, Rafieian-Kopaei M. Ethanolic Extract of Garlic for Attenuation of Gentamicin-induced Nephrotoxicity in Wistar Rats. *Iran J Kidney Dis.* 2013 Sep;7(5):376-82.
 61. Rafieian-kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P. Effects of *Ferulago angulata* Extract on Serum Lipids and Lipid Peroxidation. *Evid-Based Compl Alt.* 2014.
 62. Rabiei Z, Rafieian-kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *Zizyphus jujube* Extract on Memory and Learning Impairment Induced by Bilateral Electric Lesions of the Nucleus Basalis of Meynert in Rat. *Neurochem Res.* 2014 Feb; 39(2):353-60.
 63. Jafarzadeh L, Rafieian-Kopaei M, Samani RA, Asgari A. The Effect of Hydroalcoholic Extract of *Stachys Lavandulifolia* Vahl on Pregnant Mice. *Excli J.* 2012;11:357-62.
 64. Saki K, Bahmani M, Rafieian-Kopaei M. The effect of most important medicinal plants on two important psychiatric disorders (anxiety and depression)-a review. *Asian Pac J Trop Med.* 2014 Sep;7:S34-S42.
 65. Rabiei Z, Rafieian-Kopaei M, Mokhtari S, Shahrani M. Effect of Dietary Ethanolic Extract of *Lavandula officinalis* on Serum Lipids Profile in Rats. *Iran J Pharm Res.* 2014;13(4):1295-301.
 66. Asadi SY, Parsaei P, Karimi M, et al. Effect of green tea (*Camellia sinensis*) extract on healing process of surgical wounds in rat. *Int J Surg.* 2013; 11: 332-7.
 67. Gupta A, Shaikh AC, Chaphalkar SR. Aqueous extract of *Calamus rotang* as a novel immunoadjuvant enhances both humoral and cell mediated immune response. *J Herbmed Pharmacol.* 2017;6(1):43-48.
 68. Vahedi G, Khosravi AR, Shokri H, Moosavi Z, Delirez N, Sharifzadeh A, et al. Fungicidal effect of *Origanum vulgare* essential oil against *Candida glabrata* and its cytotoxicity against macrophages. *J Herbmed Pharmacol.* 2016;5(2):78-84.
 69. Sedighi M, Rafieian-Kopaei M, Noori-Ahmadabadi M, Godarzi I, Baradaran A. In Vitro impact of hydro alcoholic extract of *Rosa damascena* Mill. on rat ileum contractions and the mechanisms involved. *International Journal of Preventive Medicine.* 2014 ; 5(6):767-75.
 70. Sedighi M, Noori-Ahmadabadi M, Rafieian-Kopaei M, Ebrahimipoor-Samani J, Shahinfard N. The effect of *Rosa damascena* Mill hydro-alcoholic extract on the ileum contraction in rat. *J Mazand Univ Med Sci.* 2014; 23(108):30-9.
 71. Rabiei Z, Rafieian-Kopaei M. Neuroprotective effect of pretreatment with *Lavandula officinalis* ethanolic extract on blood-brain barrier permeability in a rat

- stroke model. *Asian Pac J Trop Med.* 2014 Sep;7:S421-S6.
72. Fathi F, Oryan S, Rafieian-Kopaei M, Eidi A. Neuroprotective effect of pretreatment with menthalongifolia l. extract on brain ischemia in the rat stroke model. *Arch Biol Sci.* 2015;67(4):1151-63.
 73. Nasri H, Rafieian-Kopaei M. Metformin and Diabetic Kidney Disease: A Mini-Review on Recent Findings. *Iran J Pediatr.* 2014 Oct;24(5):565-8.
 74. Nasri H, Rafieian-Kopaei M. Metformin: Current knowledge. *J Res Med Sci.* 2014 Jul;19(7):658-64.
 75. Sharafati-Chaleshtori R, Rokni N, Rafieian-Kopaei M, Drees F, Sharafati-Chaleshtori A, Salehi E. Use of Tarragon (*Artemisia Dracunculus*) Essential Oil as a Natural Preservative in Beef Burger. *Ital J Food Sci.* 2014;26(4):427-32.
 76. Bahmani M, Shirzad H, Majlesi M, Shahinfard N, Rafieian-Kopaei M. A review study on analgesic applications of Iranian medicinal plants. *Asian Pac J Trop Med.* 2014 Sep;7:S43-S53.
 77. Namjoo A, Nasri H, Talebi-Juneghani A, Baradaran A, Rafieian-Kopaei M. Safety Profile of *CarthamusTinctorius L.* in Lactation: Brain, Renal and Hepatotoxicity. *Pak J Med Sci.* 2013 Jan-Mar;29:378-83.
 78. Hasanpour Z, Nasri H, Rafieian-Kopaei M, Ahmadi A, Baradaran A, Nasri P, et al. Paradoxical Effects of Atorvastatin on Renal Tubular Cells An Experimental Investigation. *Iran J Kidney Dis.* 2015 May;9(3):215-20.
 79. Rouhi-Boroujeni H, Heidarian E, Rouhi-Boroujeni H, Deris F, Rafieian-Kopaei M. Medicinal Plants with multiple effects on cardiovascular diseases: a systematic review. *Curr Pharm Des.* 2017; 23(7): 999 – 1015
 80. Sarrafchi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative stress and Parkinson's disease: New hopes in treatment with herbal antioxidants. *Curr Pharm Des.* 2016; 22(2): 238 – 246.
 81. Shayganni E, Bahmani M., Asgary S, Rafieian-Kopaei M. Inflammaging and cardiovascular disease: Management by medicinal plants, *Phytomedicine.* 2015, <http://dx.doi.org/10.1016/j.phymed.2015.11.004>
 82. Rabie Z, Gholami M, Rafieian-Kopaei M. Antidepressant effects of *Menthapulegium* in mice. *Bangladesh J Pharmacol.* 2016; 11(3): 711-715. doi: <http://dx.doi.org/10.3329/bjp.v11i3.27318>.
 83. Sharafati-Chaleshtori R, Shirzad H, Rafieian-Kopaei M, Soltani A. Melatonin and human mitochondrial diseases. *J Res Med Sci* 2016;21:138.
 84. Rafieian-Kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P. Effects of *Ferulagoangulata* extract on serum lipids and lipid peroxidation. Evidence-Based Complementary and Alternative Medicine; 2014 (2014), Article ID 680856, 4 pages <http://dx.doi.org/10.1155/2014/680856>
 85. Bahmani M, Sarrafchi A, Shirzad H, Rafieian-Kopaei M. Autism: Pathophysiology and promising herbal remedies. *Curr Pharm Des.* 2016; 22(3):277–285.
 86. Rafieian-Kopaei M, Nasri H, Baradaran A. *Teucriumpolium*: Liver and kidney effects. *J Res Med Sci.* 2014 May;19(5):478-9.
 87. Nasri H, Rafieian-Kopaei M. Tubular Kidney Protection by Antioxidants. *Iran J Public Health.* 2013 Oct;42(10):1194-6.
 88. SalarAmoli J, Sadighara P, BarinA, Yazdani A, Satari S. Biological screening of *Amaranthusretroflexus L.* (Amaranthaceae). *Brazilian Journal of Pharmacognosy* 2009; 19(2B): 617-620.