

Research Article

Frequency of Diastolic Dysfunction in Cases of Diabetes Mellitus

¹Mehwish Khalid, ²Hurria Waheed
and ³Attia Riaz

¹Services Hospital, Lahore

²Jinnah Hospital, Lahore

³DHQ Hospital Jhang

ABSTRACT

Objective; Frequency of diastolic dysfunction in cases of Diabetes Mellitus.

Methodology; This cross sectional study was conducted during July 2017 to December 2017 at Services hospital, Lahore. Hundred cases of diabetes mellitus of both gender i.e. male/female with age more than 40 years and duration of DM of at least 2 years and of type II DM were included. The cases with previous history of acute coronary syndrome, valvular disease, liver or renal disease were excluded. The cases were labelled as Diastolic dysfunction where E/A ratio was <0.8 assessed on transthoracic echocardiography.

Results: In this study, out of 100 cases there were 61 (61%) males and 39 (39%) females with mean age of 54.34±9.41 years. There were 36 (36%) cases had HTN. Diastolic dysfunction was observed in 49 (49%) of cases. There was no significant difference in terms of gender in both groups with p= 0.79. Diastolic dysfunction was more in cases that had HTN where it was seen in 25 (69.44%) of cases with p= 0.001. There was significant difference in cases that had BMI more than 30 where it was seen in 30 (62.50%) of cases with p= 0.001.

Conclusion: Diastolic dysfunction is seen in almost half of the cases with DM and it is significantly high in cases that had HTN and BMI more than 30.

Keywords: Diastolic dysfunction, Diabetes Mellitus

INTRODUCTION;

Diabetes Mellitus (DM) is a syndrome of clinical symptoms, who is widely prevalence all across the globe and is considered as one of the most important and prevalent controllable entity. It can be classified into type I and type II where the former is present at childhood and the latter one develops at later ages. Insulin deficiency and resistance are the two major types to lead to its development.¹

DM can lead to wide range of complications as it is a systemic syndrome and can virtually involve any organ of the body. The side effects are usually sub classified into micro and macro vascular complications for which there is complex underlying pathophysiology. Chronic hyperglycemic level and formation of

atherosclerotic plaques are common modalities that are faced.²⁻³

Heart failure especially diastolic dysfunction is one of the major concern in recent times that is being focused as compared to systolic dysfunctions. According to a study more than 70% of the cases with diabetes mellitus due to a cardiovascular disease. Electrocardiogram is the initial and the echocardiography is the investigation of choice to label diastolic dysfunction.⁴⁻⁵

Multiple studies have been done in the recent past to look for this functions and have revealed its incidence from 30 to 80% of the cases. Sharavanan TKV et al, in their study found this prevalence in 55% of cases.⁶ Dikshit et al and

Srifevi et al showed that the diastolic dysfunction was seen in 66-79% of the cases in their studies respectively.⁵

Objective; Frequency of diastolic dysfunction in cases of Diabetes Mellitus.

MATERIALS AND METHODS:

Study settings; Cross sectional study

Study place; Department of Cardiology, Services Hospital, Lahore.

Study duration; July 2017 to December 2017

Sampling technique; Non probability consecutive sampling Hundred cases of diabetes mellitus of both gender i.e. male/female with age more than 40 years and duration of DM of at least 2 years and of type II DM were included. The cases with previous history of acute coronary syndrome, valvular disease, liver or renal disease were excluded. The cases were labelled as Diastolic dysfunction where E/A ratio was <0.8 assessed on transthoracic echocardiography.

Figure 01. Diastolic dysfunction



Statistical analysis;

The data was assessed by the help of SPSS version 23. After stratification of confounders stratification chi square test was applied taking p value < 0.05 as significant.

RESULTS;

In this study, out of 100 cases there were 61 (61%) males and 39 (39%) females with mean age of 54.34±9.41 years. There were 36 (36%) cases had HTN. Diastolic dysfunction was observed in 49 (49%) of cases as in figure 1. There was no significant difference in terms of gender in both groups with p= 0.79 (table 01). Diastolic dysfunction was more in cases that had HTN where it was seen in 25 (69.44%) of cases with p= 0.001 as in table 2. Table 03 reveals significant difference in cases that had BMI more than 30 where it was seen in 30 (62.50%) of cases with p= 0.001.

Table 1. Diastolic dysfunction in study subjects vs Gender

| Gender | Diastolic dysfunction | | Total | p value |
|--------------|-----------------------|-----------------|-------------------|---------|
| | Yes | No | | |
| Male | 32 (52.46%) | 29 (47.54%) | 61 (100%) | 0.79 |
| Female | 17 (43.59%) | 22 (56.41%) | 39 (100%) | |
| Total | 49(49%) | 51 (51%) | 100 (100%) | |

Table 2. Diastolic dysfunction in study subjects vs HTN

| Hypertension | Diastolic dysfunction | | Total | p value |
|--------------|-----------------------|-----------------|-------------------|---------|
| | Yes | No | | |
| Yes | 25 (69.44%) | 11 (30.56%) | 36 (100%) | 0.001 |
| No | 24 (37.50%) | 40 (62.50%) | 64 (100%) | |
| Total | 49(49%) | 51 (51%) | 100 (100%) | |

Table 3. Diastolic dysfunction in study subjects vsBMI

| BMI | Diastolic dysfunction | | Total | p value |
|--------------|-----------------------|-----------------|-------------------|---------|
| | Yes | No | | |
| More than 30 | 30 (62.50%) | 18 (37.50%) | 48 (100%) | 0.001 |
| 30 or less | 19 (36.54%) | 33 (63.46%) | 52 (100%) | |
| Total | 49(49%) | 51 (51%) | 100 (100%) | |

DISCUSSION;

The incidence of Diabetes mellitus is increasing globally and so are developing its various complications due to poor control; as it is also found in high number in developing countries. Cardiac complications like diastolic dysfunction can add to over all morbidity in such cases and warrant early diagnosis and management.

Diastolic dysfunction was seen in 49 (49%) of the cases in this study with DM. This finding was close the studies done in the past. Sharanavan et al, carried out a study on cases of DM and it was seen that diastolic dysfunction was observed in 66 (55%) of cases in their study.⁶ Similar was seen in the study of Patil et al, were this was observed in 54.33% of the wit DM.⁸

There was significant difference of DM with Diastolic dysfunction with respect to BMI more than 30 where it was seen in 30 (62.5%) of the cases with p values of 0.001. In the past it was well studies that obesity was found as a single predictor associated with diastolic dysfunction. Alfried et al, conducted a case control study and it was seen that this dysfunction was significantly higher in diabetes as compared to non diabetic with p <0.05.⁹⁻¹⁰ Russo, et al, looked for correlation of diastolic dysfunction and DM and it was also found to be a liner correlation.¹⁰

There was also significant difference in cases that had HTN where it was seen in 25 (69.44%) of cases with p= 0.001. The results of the studied done by Abhay Kumar et, also revealed that the

HTN revealed significant association of HTN with diastolic dysfunction with p=0.03.¹¹

CONCLUSION;

Diastolic dysfunction is seen in almost half of the cases with DM and it is significantly high in cases that had HTN and BMI more than 30.

REFERENCES;

1. Fontes-Carvalho R, Ladeiras-Lopes R, Bettencourt P, Leite-Moreira A, Azevedo A. Diastolic dysfunction in the diabetic continuum: association with insulin resistance, metabolic syndrome and type 2 diabetes. *CardiovascDiabetol.* 2015;14:4.
2. Chaudhary AK, Aneja GK, Shukla S, Razi SM. Study on diastolic dysfunction in newly diagnosed Type 2 diabetes mellitus and its correlation with glycosylated haemoglobin (HbA1C). *J ClinDiag Res.* 2015;9(8):OC20-OC22.
3. Patil MB, Burji NP. Echocardiographic evaluation of diastolic dysfunction in asymptomatic type 2 diabetes mellitus. *J Assoc Physicians Ind.* 2012;60:23-26.
4. Rubler S, Deluges J, Yuceoglu YZ, Kumar T, Branwood AW, Grishman A. New type of cardiomyopathy associated with diabetic glomerulosclerosis. *Am J Cardiol.* 1972;30:595-602.
5. Sridevi, Jain R, Patil R. Assessment of left ventricular diastolic dysfunction by doppler echocardiography in patients of diabetes

- mellitus. *Ind J Basic Applied Med Res.* 2015;4(2):568-70.
6. Sharavanan TKV, Prasanna KB, Ekanthalingam S, Sundaram A, Premalatha E, Arumugam B. A study on the prevalence of diastolic dysfunction in type 2 diabetes mellitus in a tertiary care hospital. *Int Arch Integ Med.* 2016;3(7):216-21.
 7. Dikshit NM, Wadia PZ, Shukla DK. Diastolic dysfunction in diabetes mellitus. *Nat J Med Res.* 2013;3(3):249-52.
 8. VirendraC.Patil, HarshaV.Patil, Kuldeep B. Shah, Jay D. Vasani, Pruthvi Shetty. Diastolic dysfunction in asymptomatic type 2 diabetes mellitus with normal systolic function. *J Cardiovasc Dis Res,* 2011;2(4):213–222.
 9. Germing A, Gotzmann M, Schikowski S, Vierkotter A. Diastolic dysfunction without abnormalities in left atrial and left ventricular geometry does not affect quality of life in elderly women. *ExpClinCardiol,* 2011;16(2):37–39.
 10. Russo C, Jin Z, Homma S, Rundek T, Elkind MS, Sacco RL et al. Effect of obesity and overweight on left ventricular diastolic function: a community-based study in an elderly cohort. *J Am CollCardiol.,* 2011; 57:1368-76.
 11. Kumar A, Girish Kumar, Aneja, ShubhraShukla, Syed MohdRazi. Study On Diastolic Dysfunction in Newly Diagnosed Type 2 Diabetes Mellitus and its Correlation with Glycosylated Haemoglobin (HbA1C). *Journal of Clinical and Diagnostic Research,* 2015;9(8): OC20-OC22.