

Research Article**Assessment of mode of retroperitoneal organ injuries
following blunt abdominal trauma****¹Muhammad Javeed Ali ²Sara Sardar
and Muhammad Yasir iqbal**¹Medical Officer, DHQ Hospital Muzaffargarh
²Woman Medical Officer,
Rural Health Centre Dullewala, Bhakkar**ABSTRACT****Objective:** To find out the frequency of mode of retroperitoneal organ injuries following blunt abdominal trauma.**Material and methods:** This cross sectional study was conducted at DHQ Hospital, DHQ Hospital Muzaffargarh from March 2017 to September 2017. Total 100 cases of abdominal trauma was selected.**Results:** Total 100 cases having history of blunt trauma abdomen were selected for the present study and average age of cases was 38.12±11.34 years. In 61 (61%) cases mode of injury was road accident followed by physical assault 11 (11%) and fall from height 28 (28%). Pancreas injury was seen in 24 (24%) cases, injury of Duodenum was noted in 28 (28%) and kidney was injured in 32 (32%) cases. In significant association of mode of injury with gender and age of the cases were noted. **Conclusion:** Most frequent reason of blunt abdomen trauma was road accidents and most of the cases were found with injury of kidney.**Keywords:** Retroperitoneal organs, blunt trauma, Road traffic accidents.**INTRODUCTION**

Like other developing countries, trauma is one of the leading preventable cause of mortality and morbidity in Pakistan.¹ In Pakistan accidental trauma ranked four among the chief causes of death. Trauma accounts about 8% of all the deaths in our country.

Approximately, 140,000 individuals die in accidents, and approximately double the number are disabled by trauma yearly.²

Trauma is defined as damage to the body by exchange with environmental energy that is beyond body's resilience.^{3,4} Due to large surface area, abdomen is the most frequently injured regions of the human body.⁵ The retroperitoneum is that part of abdomen which is separated from peritoneum anteriorly by the posterior peritoneal

fascia and is bounded posteriorly by the fascia transversalis. It contains portions of the duodenum and colonas well as the kidneys, adrenal glands, pancreas, inferior vena cava (IVC) and abdominal aorta. About 75% of abdominal trauma follows blunt injury.⁶ Associated injuries of blunt abdominal trauma are head injury, chest trauma and bony injury. Moreover, the decision to perform laparotomy for blunt abdominal trauma is more complex and difficult, as structural injury being less obvious.

The retroperitoneum is one of the most challenging areas of the abdomen.⁷ Injuries of the retroperitoneal organs occur mainly in patients with polytrauma.⁸ Retroperitoneal injuries are among the most lethal injuries sustained by

trauma patients and the most common modes of injury are road accidents, physical assault, fall from height and animal hits.⁹ Retroperitoneal organ injuries are known to occur in a significant minority of blunt abdominal trauma cases.¹⁰

MATERIAL AND METHODS:

This cross sectional study was conducted at DHQ Hospital, DHQ Hospital Muzaffargarh from March 2017 to September 2017. Total 110 consecutive cases sustaining blunt trauma abdomen were recruited. Before commencing the research, an approval was obtained from review committee of institution and written informed consent was taken from every patient.

All patients either male or female having age 12 to 60 years, presenting with history of blunt trauma abdomen and undergoing exploratory laparotomy were included in the study. The diagnosis of blunt abdominal trauma was made on the basis of presence of tenderness, rigidity, and bruise on the abdominal wall. Patients presenting within 12 hours of sustaining injury will be included in the study. Patients managed non-operatively, patients suffering from any kind of penetrating abdominal injury and moribund patients of ASA-5 were excluded from the study.

Demographic profile of all the subjects was entered in pre-designed Performa. Mode of injury and intra-abdominal injuries involving the kidney, pancreas and duodenum were noted. The scale devised by the Organ Injury Scaling Committee of the American Association for the Surgery of Trauma was used to grade the injuries to various organs. Grading of injuries was verified by attending consultant.

Data was analyzed by using SPSS version 17. Numerical variables were presented as mean and SD. Frequencies and percentages were calculated for categorical data. To see the association between different variable, chi square test was used and p value 5% was considered as significant.

RESULTS:

In this study, total 100 case of blunt abdomen trauma were recruited. Mean age of the cases was 38.12 ± 11.34 years. In 61 (61%) cases mode of injury was road accident followed by physical assault 11 (11%) and fall from height 28 (28%). (Fig. 1)

Pancreas injury was seen in 24 (24%) cases in which 5 (20.33%) cases found with grade I injury followed by 9 (38.37%) grade II, 5 (20.83%) grade III, 4 (16.67%) grade IV and 1 (4.17%) found with grade V injury. Injury of Duodenum was noted in 28 (28%) cases in which 4 (14.29%), 14 (50%), 6 (21.43%), 3 (10.71%) and 1 (3.57%) cases were found with Grade I, II, III, IV and V respectively. Kidney was injured in 32 (32%) cases and pancreatoduodenal injury was noted in 16 (16%) cases. (Table 1).

Out of 61 (61%) cases injured by road accident, male cases were 49 (81.82%) and female cases were 12 (19.67%). Out of 28 (28%) cases injured due to fall from height, 18 (64.29%) cases were male and 10 (35.71%) cases were female and out of 11 (11%) cases injured by physical assault, 7 (63.64%) cases were male and female cases were 4 (36.36%). Association of mode of injury with gender was not statistically significant ($P = 0.196$). (Table 2). Out of 61 cases of road accident, 41 (67.21%), 18 (29.51%) and 2 (3.28%) cases belonged to age group 12-35 years, 36-50 years and 51-70 years respectively. Out of 28 cases injured with fall from height, 14 (50%) belonged to age group 12-35 years, 9 (32.14%) belonged to age group 26-50 years and 5 (17.86%) belonged to age group 51 – 70 years.

Cases having history of physical assault was 11, of which 6 (54.55%) belonged to age group 12-35 years, 3 (27.27%) to age group 36-50 years and 2 (18.18%) cases belonged to age group 51-70 years.

But mode of injury insignificantly ($P = 0.148$) associated with age of the cases. (Table 3)

Fig. 1: Mode of injury




Table 1: Organ injuries and grades

Organ injury	Grade of injury					Total (%)
	I (%)	II (%)	III (%)	IV (%)	V (%)	
Kidney	5 (15.63)	10 (31.25)	8 (25)	6 (18.75)	3 (9.38)	32 (32)
Duodenum	4 (14.29)	14 (50)	6 (21.43)	3 (10.71)	1 (3.57)	28 (28)
Pancreas	5 (20.33)	9 (38.37)	5 (20.83)	4 (16.67)	1 (4.17)	24 (24)
Pancreatoduodenal	11 (68.75)	4 (25)	1 (6.25)	0	0	16 (16)
Total	25 (25)	37 (37)	20 (20)	13 (13)	5 (5)	100

Table 2: Gender distribution

Mode of injury	Gender		Total (%)	P. value
	Male (%)	Female (%)		
Road accidents	49 (81.82)	12 (19.67)	61 (61)	0.196
Fall from height	18 (64.29)	10 (35.71)	28 (28)	
Physical assault	7 (63.64)	4 (36.36)	11 (11)	
Total	74 (74)	26 (26)	100	

Table 3: Age distribution

Mode of injury	Age Group			Total (%)	P. value
	12-35 years	36-50 years	51-70 years		
Road Accident	41 (67.21)	18 (29.51)	2 (3.28)	61 (61)	0.148
Fall from	14 (50)	9 (32.14)	5 (17.86)	28 (28)	

Height				
Physical Assault	6 (54.55)	3 (27.27)	2 (18.18)	11 (11)
Total	61 (61)	30 (31)	9 (9)	100

DISCUSSION

In present study, male cases were more victim of abdominal trauma as compare to female cases (74% vs 26%) similar finding was reported by Khan et al,⁶ according to this study young males who belonged to age group 20 to 30 years were the most frequent victim of blunt abdominal trauma. In this study, road accident was the most common (61%) cause of blunt abdominal trauma. The second most common (28%) cause was falling from a height and the third was Physical assault (11%). Some other studies also reported that road accidents, interpersonal violence and falls from height as main causes of blunt abdominal trauma.^{12,13} Ahmed et al also reported trauma as the leading cause of mortality in subjects having age 1-44 years.⁹ Blunt abdominal trauma accounted for 79% cases and males are more victim of blunt abdominal trauma as compare to female. In one study by Bhattacharjee et al,¹⁴ blunt abdominal trauma is more frequent in males aged 21-30 years; the majority of patients were injured in automobile accidents.

In our study the most common injured organ was kidney (32%) and second most common (28%) injured organ was duodenum. A study by Zaydfudim et al.,¹⁵ indicated that most of patients with vertical deceleration injuries (i.e., falls from heights), only 5.9% had blunt abdominal injuries. Consistent with these results, studies by Bhattacharjee et al¹⁴ and Antonacci et al,¹⁶ injuries to the duodenum account for approximately 3% to 5% of abdominal trauma. Blunt abdominal trauma as a result of direct blow to the epigastrium, mainly due to road traffic accident and sports trauma (bicycle handle injury), accounts for 25% of all duodenal injuries as shown by Chinnery et al and Girgin et al.^{17,18}

In our study kidney damage was noted in 32% cases. Grad I injury was seen in 15.63% cases, Grade II, III, IV and V injuries were seen 31.25%, 25%, 18.75% and 9.38% respectively. Similar

findings were reported by Wong et al,¹⁹ 89 cases of Grade 2 renal injuries were recorded with blunt trauma accounting for 94.4%; 57.3% were Grade 2 injuries, 12.4% Grade 3, 25.8% Grade 4, and 4.5% Grade 5. MVAs and motorcycle accidents were the most common cause of injury, accounting for 48.3% of all renal injuries.

In our study pancreas was injured in 24% cases, whereas 16% had pancreato-duodenal injuries. Traumatic injuries of the pancreas occur after blunt abdominal traumas or penetrating wounds with a ratio of 3:1.¹⁶ These are characterized by high morbidity and mortality with a 45-50% combined rate as reported in the reviewed literature.^{17,20} Pancreatic injuries occur in 3-15% of all abdominal trauma. Isolated traumatic injuries of the pancreas are uncommon; in 50-98% of cases they are associated with injuries to other organs, such as spleen, liver, kidney, large/small intestine, veins or arteries. Due to the retroperitoneal location of the pancreas, isolated pancreatic injury occurs in less than 5% of cases of major blunt abdominal trauma.²¹

CONCLUSION:

Road accidents were the most common cause of blunt abdominal trauma and most of the cases were found with injury of kidney. This study highlights the need for prioritizing a public health approach to abide by traffic laws and violence prevention in Pakistan.

REFERENCES:

1. Jat AA, Khan MR, Zafar H, Raja AJ, Hoda Q, Rehmani R, et al. Peer review audit of trauma deaths in a developing country. *Asian J Surg.* 2004 Jan;27(1):58-64.
2. Nasrullah M, Muazzam S. Risk differences between children and adults in road traffic injuries: a descriptive study from a tertiary-care hospital in a low-income country. *Eur J Emerg Med.* 2012 Jun;19(3):167-70.

3. Bhan C, Forshaw MJ, Bew DP, Kapadia YK. Diagnostic peritoneal lavage and ultrasonography for blunt abdominal trauma: attitudes and training of current general surgical trainees. *Eur J Emerg Med.* 2007 Aug;14(4):212–5.
4. MacLeod JBA, Cohn SM, Johnson EW, McKenney MG. Trauma deaths in the first hour: are they all unsalvageable injuries? *Am J Surg.* 2007 Feb;193(2):195–9.
5. Kuncir EJ, Velmahos GC. Diagnostic peritoneal aspiration--the foster child of DPL: a prospective observational study. *Int J Surg.* 2007 Jun;5(3):167–71.
6. Khan JS, Iqbal N, Gardezi JR. Pattern of visceral injuries following blunt abdominal trauma in motor vehicular accidents. *J Coll Physicians Surg Pak.* 2006 Oct;16(10):645–7.
7. Naguib NNN. Delayed presentation of shock due to retroperitoneal hemorrhage following a fall. *J Emerg Trauma Shock.* 2009;2(2):139–43.
8. Sica G, Bocchini G, Guida F, Tanga M, Guaglione M, Scaglione M. Multidetector computed tomography in the diagnosis and management of renal trauma. *Radiol Med.* 2010 Sep;115(6):936–49.
9. Ahmed N, Vernick JJ. Pancreatic injury. *South Med J.* 2009 Dec;102(12):1253–6.
10. Daly KP, Ho CP, Persson DL, Gay SB. Traumatic Retroperitoneal Injuries: Review of Multidetector CT Findings. *RadioGraphics.* 2008 Oct;28(6):1571–90.
11. Mofidi M, Hasani A, Kianmehr N. Determining the accuracy of base deficit in diagnosis of intra-abdominal injury in patients with blunt abdominal trauma. *Am J Emerg Med.* 2010 Oct;28(8):933–6.
12. Terreros A, Zimmerman S. Duodenal hematoma from a fall down the stairs. *J Trauma Nurs.* 2009 Sep;16(3):166–8.
13. Khan AR, Fatima N, Anwar K. Pattern and management of renal injuries at Pakistan Institute of Medical Sciences. *J Coll Physicians Surg Pak.* 2010 Mar;20(3):194–7.
14. Bhattacharjee HK, Misra MC, Kumar S, Bansal VK. Duodenal perforation following blunt abdominal trauma. *J Emerg Trauma Shock.* 2011;4(4):514–7.
15. Zaydfudim V, Cotton BA, Kim BD. Pancreatic transection after a sports injury. *J Trauma.* 2010 Oct;69(4):E33.
16. Antonacci N, Di Saverio S, Ciaroni V, Biscardi A, Giugni A, Cancellieri F, et al. Prognosis and treatment of pancreaticoduodenal traumatic injuries: which factors are predictors of outcome? *J Hepatobiliary Pancreat Sci.* 2011 Mar;18(2):195–201.
17. Chinnery GE, Madiba TE. Pancreaticoduodenal injuries: re-evaluating current management approaches. *S Afr J Surg.* 2010 Feb;48(1):10–4.
18. Girgin S, Gedik E, Yağmur Y, Uysal E, Baç B. Management of duodenal injury: our experience and the value of tube duodenostomy. *Ulus Travma Acil Cerrahi Derg.* 2009 Sep;15(5):467–72.
19. Wong KY, Brennan JA, Calvert RC. Management of severe blunt renal trauma in adult patients: a 10-year retrospective review from an emergency hospital. *BJU Int.* 2012 Oct;110(8):E330.
20. Shoobridge JJ, Corcoran NM, Martin KA, Koukounaras J, Royce PL, Bultitude MF. Contemporary Management of Renal Trauma. *Rev Urol.* 2011;13(2):65–72.
21. Ankouz A, Elbouhadouti H, Lamrani J, Taleb KA, Louchi A. Pancreatic transection due to blunt trauma. *J Emerg Trauma Shock.* 2010;3(1):76–8.